

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
13 January 2005 (13.01.2005)

PCT

(10) International Publication Number
WO 2005/003364 A2

(51) International Patent Classification⁷: C12N 15/90

(21) International Application Number:
PCT/GB2004/002869

(22) International Filing Date: 1 July 2004 (01.07.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0315379.8 1 July 2003 (01.07.2003) GB

(71) Applicant (for all designated States except US): **OXITEC LIMITED** [GB/GB]; 71 Milton Park, Abingdon, Oxfordshire OX14 4RX (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **ALPHEY, Luke** [GB/GB]; 71 Milton Park, Abingdon, Oxfordshire OX14 4RX (GB).

(74) Agent: **HILTON, David**; Marks & Clerk, 57-60 Lincoln's Inn Fields, London WC2A 3LS (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: STABLE INTEGRANDS

(57) Abstract: There is provided a transposable element comprising at least four inverted repeats, at least two of which are each inverted in relation to another, the element comprising DNA for insertion into a host genome, the DNA being located between two pairs of opposing repeats excisable by a transposase *in situ* to leave said DNA without flanking transposon-derived repeats in the host genome. Also provided is a transposable element comprising at least three inverted repeats, at least one of which is inverted in relation to the others, wherein at least one non-terminal repeat is a minimal repeat. Both these elements allow for greater efficiency of insertion of nucleotide sequences into the genome.



WO 2005/003364 A2

STABLE INTEGRANDS

The present invention relates to transposable elements and methods for their incorporation into the genome.

Transposable element technology has enabled the genetic transformation of a wide range of insects [*c.f.* Handler, A. (2001), *Insect Biochem Mol. Biol.* 31, 111-128; Handler, A. (2002), *Insect Biochem. Mol. Biol.* 32, 1211-1220; Horn, *et al.*, (2002), *Insect Biochem. Mol. Biol.* 32, 1221-1235], and at least some of these transposable elements have also been shown to be mobile over a much wider phylogenetic range, including prokaryotes and vertebrates (Handler, 2001, *supra*).

Transposons have been described extensively in the prior art. The elements used for genetic transformation of insects are generally characterised by opposing inverted repeat sequences and are associated with an optionally integral transposase enzyme. The transposase recognises the inverted repeat sequence and excises the sequences, together with any intervening DNA, and then reinserts the resulting transposon at another site, either in the genome or in a plasmid.

An autonomous transposon encodes its own transposase; for a type II transposon this is in the DNA between the inverted repeat sequences. Such autonomous transposons are not stably incorporated into the genome, as they are liable to move at random. Accordingly, where such a transposon is to be used to incorporate target DNA into a genome, it is preferred to provide the transposase on a helper plasmid, for example, so that transposition is dependent on the availability of the helper plasmid.

This system for incorporating genes or nucleic acid sequences into genomes is effective, but the transposed DNA is still associated with the inverted repeat sequences of the transposon, so that any future exposure to transposase can result in movement of the transposon, possibly even across the species barrier, although this unlikely.

The presence, or potential presence, of transposases capable of remobilising these insertions has led to objections to the use of this technology, particularly when the transformed organisms are for release into the environment. In addition, the presence of such transposases

will tend to destabilise the insertions, which is generally undesirable. Though laboratory strains can be screened for the presence of related transposases by any of several methods, based, for example, on nucleic acid sequence similarity, such as by hybridisation or PCR, or function, such as plasmid-to-plasmid transposition assays or plasmid excision assays, it is not possible exhaustively to test all wild populations.

It is therefore desirable to develop transformation methods which lead to a transgenic line which is insensitive to transposase, or in which the insertion is insensitive to transposase.

In order to stabilise the genetic insert, it is possible to provide three or more repeats, and locate the insert between two repeats in the same orientation. It follows that the number of transposable elements is equivalent to the number of repeats in one orientation multiplied by the number of repeats having the opposite orientation. Exposing the construct to transposase will result in all of the possible transposable elements being obtained, including the entire length of the transposon containing the gene of interest, which can then be inserted into the genome. The resulting insertion can then lose the shorter length which does not contain the insert by further, or the same, transposase activity. This shorter length is a transposable element flanked by inverted repeats. Once removed, it leaves the desired genetic insert in place, but with only one repeat i.e. no longer within a functional transposable element, thereby disabling the ability of transposase to excise the genetic insert. This process is subject to extremely low success rates, however, as the transposase will usually generate the shortest length transposable element.

Thus, transposable elements with repeated ends can potentially utilise any of their repeated ends for transposition or excision. It is therefore possible to obtain transposition using a first distal end, followed by excision using another, more proximal end, leaving a fragment behind. This scheme is illustrated in Figure 1.

In Figure 1, triangles A-C represent the functional ends of the transposon, for example, the short inverted repeats at the ends of a class II element such as *piggyBac*. Transposition can occur between any two opposed triangles: in this case, A-B and A-C. Flanking transposon DNA may also be included, see below. An appropriate target site sequence, for example, TTAA or (A/T)N(A/T)TTAA(A/T)N(A/T) is also assumed to be included in these examples. Proximal and distal are relative to the 5' end. The construct of Figure 1A is introduced into cells or embryos, for example, by microinjection, transfection, or ballistic or other methods, with a

suitable transposase helper, for example, helper plasmid, RNA, transposase protein or integrated transposase source, also known as a jumpstarter.

A transformant containing the integrated product must then be selected, as shown in Figure 1B.

The selected transformant is then exposed to a suitable source of active transposase, for example, helper plasmid, RNA, transposase protein or an integrated transposase source, to generate the specific excision product shown in Figure 1C. The resulting product is substantially or completely stable to further exposure to active transposase, relative to the starting construct.

In the above example, two integration events are possible in the initial transformation step, then two excisions are possible once the desired integrant has been obtained. The alternative integration utilises the proximal 3' end, resulting in the insertion of a smaller transposon than desired, lacking the DNA of interest; the alternative excision utilises the distal 3' end, resulting in the complete excision of the construct and hence excision of the DNA of interest. Insect transformation is an inefficient process, so it is desirable to optimise the efficiency of this step. With suitable markers, even inefficient or rare excision may be acceptable, as an integrated transposase source can be used, so that it is possible to score large numbers of insects without requiring microinjection.

It is, therefore, highly desirable to maximise the efficiency of the first step, so as to provide the greatest possible number of initial transformants obtained by microinjection, for example, and obtain as many full-length insertions as possible and to maximise the ratio of full-length to short insertions.

The "3-end" system described above and illustrated in Fig.1 has another weakness: the presence in the final product of a transposon end adjacent to the DNA of interest. In principle, insertion of another identical or cross-mobilising transposable element nearby could reconstitute a composite transposon and thereby destabilise the DNA of interest. There is, therefore, a need for a method to generate insertions in which no transposon ends remain.

Another available option for insect transformation involves adding another mobile element near or at one end of the transposon and rely on imprecise excision of this element to

stabilise the insertion by deleting one end. This is an inherently unsatisfactory method as the nature of the deletion is unpredictable and uncontrollable.

Another method for generating transgenics without flanking transposon DNA is well known in *Drosophila*, though not for any other insect. This is the homologous recombination system of Rong and Golic (Rong, Y., and Golic, K. (2000), *Science* 288, 2013-2018; Rong, Y., and Golic, K. (2001), *Genetics* 157, 1307-1312). Essentially, the contents of a transposon insert are moved to another, predetermined chromosomal location. The new insertion need not have any transposon DNA, though it does have to have a target site for a site-specific recombinase, such as Flp/FRT. As originally described, this insertion is always associated with a large direct repeat. The main limitation of this scheme is the highly variable, and generally very low, rate at which the new homology-based insertions are recovered. It also requires several enzyme activities to be available, such as by prior establishment of transgenic lines, for example FLP and I-SceI. The main use of this system is to generate knock-outs or other targeted modifications to specific genes or sequences.

Various attempts in the art have been made to improve the effectiveness of transposition. WO 02/46444 discloses a transposable polynucleotide suitable for use in methods for manipulating nucleic acids to create libraries of cells containing transposed nucleic acid, wherein the transposable polynucleotide comprises two or more inverted repeat sequence pairs. Each pair has a distinct and separable ability to interact with a distinct transposase enzyme. The pairs can, for instance, be provided in a nested fashion such that both members of one pair are flanked by both members of the second pair. For instance, the Tn5 transposon, described therein, includes "inside end" sequences and "outside end" sequences. A transposase is disclosed that preferentially binds and interacts with the outside ends but not the inside ends.

WO 01/91802 discloses a chimeric nucleic acid vector comprising a denoviral inverted terminal repeat flanking regions comprising, between the inverted terminal repeats, retroviral long terminal repeat flanking regions, within which is further found a cassette comprising the DNA of interest and a *gag* nucleic acid region. Such vectors are capable of transducing all *cis* and *trans* components of a retroviral vector for the generation of a high titer recombinant retroviral vector for use in *in vivo* gene transfer applications.

Wobus *et al.*, Molecular and General Genetics, 1990, vol. 222, pp 311-316 describes a new transposable element from *Chironomus thummi* which is shown to have inverted repeats of 17 base pairs at each termini. The transposable element is called TECth1, is 1.7kb long and was found in the 3' flanking region of a *C. thummi* Balbiani ring gene.

US 2002/0173634 sets out to solve problems in the use of the piggyBac vector caused by lack of suitable restriction sites to cut the components needed for gene transfer, and limitations on the sizes of genes transferred by use of this vector. This is achieved by removing large portions of apparently non-essential DNA from the piggyBac transformation vector.

US 6,200,800 B1 discloses retroviral vectors containing *cis*-acting viral elements from the expression, encapsidation, reverse transcription and integration of the retroviral genome nucleic acid sequence. A retroviral vector is provided for eliminating most of the viral elements which are not useful in the integrated provirus. The vector provided use, among other thing, the bacteriophage P1 Cre-lox recombination system.

Russ *et al.*, J. Virology, 1996, vol. 70, pp. 4927-4932 discloses self-deleting retrovirus vectors for use in gene therapy. Russ *et al.* discloses how retrovirus vectors containing a *loxP* site fuse to independently expressed selectable marker generates proviruses flanked by loxP. This enables the Cre recombinase to excise most of the provirus apart from the marker gene.

Steiner S. *et al.*, Genetics, 1995, vol. 140, pp. 973-987 shows that homologous recombination is the main mechanism for DNA integration and the cause of rearrangements in the filamentous ascomycete *Ashbya gossypii*.

What is required is a simple system with enhanced rates of initial transformation with a desired gene or nucleotide sequence.

Thus, according to a first aspect, the present invention provides a transposable element comprising at least four inverted repeats, forming at least two pairs of opposing pairs of inverted repeats, the element comprising DNA for insertion into a host genome, the DNA being located between two pairs of opposing repeats such that excision by a transposase or transposases of said pairs, *in situ*, is effective to be able to leave said DNA integrated into the host genome, without the presence of flanking transposon-derived repeats derived from said transposable element.

The DNA for insertion into a host genome is preferably a gene that is to be expressed in the host, although it is envisaged that it could preferably also be a genetic element that does not encode a protein, such as regulatory element, for instance a promoter or enhancer sequence.

Alternatively, it is also preferred that the DNA for insertion into a host genome is simply a DNA sequence for insertion into a target sequence for the purposes of, for instance, altering the frame of a coding sequence, disrupting the function of a gene or even deleting a portion of DNA if, for instance, the transposition occurs between target sites some distance apart.

A gene's function may be disrupted, for instance, by insertion of stop codon into the coding sequence thereof, insertion of suitable nucleotides sufficient to bring about a stop codon where one was not previously found in frame, or insertion of "junk" DNA into the coding sequence or untranslated regions, thereby disrupting the gene's promoter or enhancer, for instance.

Preferably, the inverted repeats are "piggyBac" repeats.

It is also preferred, the number of inverted repeats is 6, 7, 8 or more, but more preferably 5 and most preferably 4, as transposases generally favour shorter sequences to transpose or excise, thus making it preferable to have as few inverted repeats as possible to minimise unwanted transpositions and excisions and to promote the intended transpositions and excisions. This is so as to, for instance, leave only the gene-of-interest remaining in the host genome, such that as much as possible of the genetic material derived from the transposon, with the exception of the gene-of-interest, is removed.

Whilst it is preferred to remove only some of the transposon-derived DNA from the host genome, such as only one excisable transposon of opposing inverted repeats, it is more preferable to excise all transposon-derived DNA from the host genome, apart from the gene or DNA of interest.

The inverted repeats, also referred to as "ends" in the art, are preferably homologous, such that the repeat sequence is the same, albeit inverted. This has the advantage of requiring only one transposase. Alternatively, it is also preferred that the different pairs of homologous

repeats are heterologous, such that they do not form opposing inverted repeats which may be transposable or excisable. Preferably, the heterologous repeats are recognised by different transposases, thereby allowing the user to exert control over different transposition and excision steps.

The transposase is preferably autonomous, such that the transposase is encoded within the transposon itself. However, it is also preferred that the transposase is provided by another means, such as a further genetic element comprising the transposase, for instance another transposon or preferably a helper plasmid, thereby also allowing the user a greater level of control as to when the various stages of the transposition and excision according to the present invention occur.

The transformant is preferably exposed to a suitable source of active transposase, for example, a helper plasmid or RNA encoding the transposase, or a transposase protein or integrated transposase source, as known in the art.

Preferably, one or more of the inverted repeats is a minimal repeat, as discussed below. In this instance, it is preferred that the minimal repeat is non-terminal, i.e. it is an internal repeat, as discussed below.

It is also preferred that markers are associated with the transposon in order to allow the user to follow the progress of the various steps of transposition and excision and to determine in which individuals have been said steps have successful. Suitable markers and systems therefor are also discussed below. Preferably at least one genetic marker is associated with an identifiable step in the transposition/excision process, and more preferably, the marker is associated with the DNA for insertion into a host genome.

Preferably, the DNA for insertion into a host genome does not comprise any inverted repeats or target sites for insertion of the present invention, which could lead to unwanted excision of part or all of a previous insertion, or to the transposable element of the present invention being transposed into a previous insertion. Although it may be impractical to have absolutely no inverted repeats or insertion sequences in the DNA for insertion into a host genome, it is preferred that these be kept to a minimum or that they are heterologous to, and therefore not recognised by, the transposase or transposases used by the present invention.

Opposing inverted repeats are capable of recognition by a suitable transposase, leading to transposition or excision of the repeat and intervening DNA. If the inverted repeats are not opposing, i.e. they are orientated in such a way that the second repeat sequence is not a complementary mirror image of the first repeat sequence, then no transposition or excision will occur between these repeats. This can be seen in Figure 1A, where repeats A and B are opposing, thereby allowing transposition/excision therebetween, whereas repeats B and C are not opposing. It can also be seen in Figure 3A, where the two repeats marked 5' are not opposing, nor is the left hand 5' repeat in relation to the left hand 3' repeat. In the diagrammatic representations used in Figures 1-4, an opposing pair of repeats is shown by a 5' arrow and a 3' pointing in opposite directions.

The opposing inverted repeats of the present invention are preferably excised *in situ*, from the host genome, following initial transposition of the full-length element, comprising the at least four repeats and the DNA of interest, into the host genome.

Preferably, the element comprises two external opposed inverted repeats, one on each side of an inversion cassette, the cassette comprising;

the DNA for insertion into a host genome, two inverted cassette repeats and two inversion sites, the DNA for insertion into a host genome being flanked on either side by one of the inverted cassette repeats, each inverted cassette repeat being further flanked by an inversion site;

the cassette being capable of inversion within the transposed element *in situ* in the presence of a recombinase, such that following inversion, the two inverted cassette repeats flanking the DNA for insertion into a host genome each separately form a further pair of opposing inverted repeats with one of the external inverted repeats, the further pairs of opposing repeats being excisable by a transposase *in situ* to leave said DNA without flanking transposon-derived repeats in the host genome.

It will, of course, be appreciated that the repeats or inversion sites that, for instance, are said to be flanking or bounding another feature, are not necessarily intended to be directly adjacent thereto, but may be proximal or have a spacer therebetween, provided that this does not impair the function of either feature. In some cases, it is envisaged that such a spacer may be useful and may even include a marker, for instance.

The external inverter repeats are found on each side of the inversion cassette. Before inversion, the external opposing repeats form a pair of opposing inverted repeats transposable by a transposase, such that the repeats and the cassette comprised therebetween are transposed into the host genome by the transposase. Following inversion of the cassette in the host genome by a recombinase, each of the external repeats then forms a new excisable element or transposon with an opposing inverted repeat that is now opposing as a result of the inversion. Two further newly-formed opposing pairs of inverted repeats are found, one on each side of the inverted cassette and are excisable. Being shorter than the pair formed between the two external repeats found at either end of the transposed and inverted element and comprising the DNA or gene of interest, the newly-formed repeats are favoured over the longer repeat and are excisable in preference thereto, leaving only the DNA of interest, preferably with minimal flanking sequences, in the host genome.

A transposon-derived repeat is a repeat that originated in the transposable element and not from the host genome, or is homologous to, complementary to or a variant of that original repeat.

The inversion sites are sequences recognised by inversion-inducing recombinase. Preferably, such sites comprise the F₁p or Cre sites from the F₁p/FRT or Cre/lox inversion systems, or any other such systems known in the art. Preferably, the inversion sites are recognised by a directional recombinase, the recombinase-mediated inversion being essentially irreversible. Preferred sites of this type are, for instance, lox66 or lox71.

The inversion cassette preferably comprises the DNA of interest, the two inverted cassette repeats which later go on to form one half of the opposing repeats that are used to excise the transposon-derived DNA, and the inversion sites, such as the FRT site of a F₁p/FRT system or a lox site from a Cre/lox system, which direct the recombinase to the inversion of the intervening DNA. Preferably, the cassette may also comprise markers, as discussed below.

The present invention also provides a method of transposition comprising use of the transposable element to insert DNA of interest into a host genome, together with the transformant or organisms produced by said method.

In one aspect the present invention preferably provides a transposable element comprising at least four inverted repeats, at least two of which are inverted in relation to the others, comprising DNA for insertion into a host genome located between two pairs of opposing repeats excisable by a transposase *in situ* to leave said DNA without flanking transposon-derived repeats in the host genome.

Inverted repeats are sequences which are found in identical (but inverted) forms. In other words, the repeats have the same sequence running in the 5' to 3' direction, except that these sequences are on different strands of the DNA. For example, 5'-CCCTAG-3' and 5'-CTAGGG-3' are inverted repeats as the second sequence is the mirror image of the first, except that the mirror image has also been converted into the antisense of the first, such that A has been converted to T, C to G and so forth. Thus, the complementary sequence of the second repeat reads 3'-GATCCC-5' (5'-CCCTAG-3'), the same of the first repeat, only on the complementary strand.

The inverted repeat sequences of piggyBac transposons, for example, are associated with short, external TTAA sequences, which generally correspond to the insertion site. It is, therefore, preferred that such sequences are provided with the terminal inverted repeats of the present invention.

Figure 3 illustrates a scheme involving two pairs of inverted repeats flanking the gene or DNA of interest, and how this may be used to generate an insert ultimately associated with no transposon repeats.

The construct of Figure 3A is injected with a suitable transposase helper, for example, helper plasmid, RNA, transposase protein or integrated transposase source.

A transformant containing an integrated product as shown in Figure 3B may then be selected, and exposed to a suitable source of active transposase, for example, helper plasmid, RNA, transposase protein or integrated transposase source.

Transformants containing either of the specific excision products shown in Figure 3C may then be selected. Further exposure to transposase may then be employed to obtain the final excision product shown in Figure 3D.

As mentioned above, the inverted repeat sequences of piggyBac transposons, for example, are associated with short, external TTAA sequences, which generally correspond to the insertion site. The inverted repeats are also associated with internal stretches of DNA. These internal stretches of DNA can be deleted to leave a minimal repeat, so that the repeats simply flank the desired DNA to be inserted. Transposons created in this manner are capable of excision and plasmid-to-plasmid transposition, and can do so with frequencies similar to those of constructs containing more internal piggyBac sequence (Elick et al., 1997, *supra*). However, such minimal piggyBac elements are not capable of efficient germline transformation, giving transposition frequencies approximately 20x lower than more complete elements.

It has also recently been established that transposable activity is retained if part of the repeat is deleted, but that the resulting insert can no longer be excised by transposase activity at the truncated repeat.

It has now been found that it is possible to provide a transposon with three or more inverted repeats of which a middle repeat is a minimal repeat, such a transposon providing significantly enhanced levels of initial transformation.

Thus, in a further aspect, the present invention provides a transposable element comprising at least three inverted repeats, at least one of which is inverted in relation to the others, wherein at least one non-terminal repeat is a minimal repeat.

Preferably, the element comprises DNA for insertion into a host genome located between the minimal repeat and a repeat having the same orientation as the minimal repeat.

The DNA for insertion into a host genome is preferably flanked by two pairs of opposing repeats excisable by a transposase *in situ* to leave said DNA without flanking repeats in the host genome, although it is preferred that each of the repeats bounding the DNA for insertion into a host genome is a minimal repeat.

It is preferred that at least one repeat distal to the DNA for insertion into a host genome in relation to a minimal repeat in the same orientation has an internal deletion or is otherwise

compromised over up to 50% of its length, thereby reducing the frequency of excision by a transposase at that repeat.

Preferably, at least one genetic marker associated with an identifiable step in the transposition/excision process and more preferably, the marker is associated with the DNA for insertion into a host genome.

It will be appreciated the greatest length of a transposon is the distance between, and including, the two opposing repeats farthest from each other. Other, shorter transposons can then be defined within the length of the longest transposon by the presence of other repeats. It is at least one of these non-terminal, or internal, repeats that are preferably minimal repeats.

As used herein, the term 'minimal repeat' applies to the highly conserved repeat sequences observed to be required for the activity of a given transposase. The piggyBac transposon, for example, has a 32bp terminal inverted repeat interrupted by a 4bp insertion at the 5' end and a 31bp insertion at the 3' end. This can be considered as two pairs of inverted repeats, one of 13bp and another of 19bp, these two being separated by 4bp at the 5' end and 31 bp at the 3' end. The minimal repeat, in this instance, then comprises a 32bp repeat with a 4 or 31 bp insertion at the 5' and 3' ends respectively.

It is generally preferred to provide the flanking sequences commonly associated with the terminal repeats. In the case of piggyBac, this is preferably TTAA.

In general, transposases will be more effective at cutting out shorter sequences so that, where a transposon has one 5' repeat and two 3' repeats, for example, the most common transposon that will be observed transferring to another locus will be the shorter, formed by the 5' repeat together with the more proximal of the two 3' repeats. This preference is both inevitable, owing to the fact that the longer transposon will still be cut by the transposase, as well as being exacerbated by topological considerations, whereby repeats in closer proximity appear to be bound substantially better by transposases.

The present invention overcomes this problem by employing at least one minimal repeat internal to the longest transposon of the construct. This minimal repeat is utilised at only very low rates for the original insertion, thereby strongly biasing the initial reaction to generating the

larger transposon. Any DNA of interest is located between the minimal repeat and a corresponding full length repeat in the same orientation.

WO 0/0441504 discloses the use of a 3 end (3 inverted approach), but there is no mention of the use of minimal ends. There is a discussion of the use of inversion, similar to that used in conjunction with the transposable element comprising at least 4 inverted repeats according to the first aspect of the present invention. However, WO 0/0441504 only discloses the use of inversion in relation to the 3 end approach and not in relation to a 4 end approach.

It will be appreciated that minimal repeats may simply comprise the minimum repeat necessary to effect transposition, and be associated with none of the original intervening DNA between the repeats observed in piggyBac, for example. However, the present invention envisages using some of the internal sequence, although it is preferred to keep this to a minimum, as greater lengths will increase the transposition frequency of the minimal repeat, thereby diminishing the desirable bias discussed above. Accordingly, while it is preferred to keep internal sequence associated with the minimal repeat to zero nucleotides, it is possible to use up to 100 bases, for example, of the original sequence, but it is preferred to use 50, or less, and preferably 10 or less.

Correspondingly, for the full length sequences, there is no clearly established limit as to what constitutes full length. It is established, however, that between about 6 and 14% of the internal sequence of a naturally occurring transposon is sufficient to provide high levels of transposition frequency, so that the present invention generally prefers that 'full length repeat sequences' be associated with at least 5% of the original internal sequence of the transposon from which they are derived, with between 6% and 14% being preferred, and 8% to 12% being more preferred.

The transposons of the present invention may employ two or more minimal repeats, although two is the preferred maximum. Where two are used, then it is preferred that these both be internal and in opposite orientations. Where two smaller transposons flank the DNA of interest, this has the particular advantage of encouraging the full length transposable element to be incorporated into the genome and then losing the two smaller transposons, thereby leaving simply the DNA of interest without any flanking repeats. This method is also provided herein, simply using full length repeats, as defined above.

It is preferred to use a minimal piggyBac end as the more proximal of the distal ends and a fully-functional piggyBac end as the distal end. This arrangement strongly biases the initial transformation step towards insertion of the desired longer transposon.

In an alternative embodiment, multiple copies of the 3' and/or 5' ends is provided, with minimal repeats preferably providing the inmost of any series of repeats in the same orientation. This increases the chance that one or more longer versions of the transposon are integrated in the initial transformation, as required. One version of this is illustrated diagrammatically in Figure 2.

The construct of Figure 2A is introduced into cells or embryos, for example, by microinjection, transfection, or ballistic or other methods, with a suitable transposase helper, for example, helper plasmid, RNA, transposase protein or an integrated transposase source.

A transformant containing the integrated product of Figure 2B is then selected, and exposed to a suitable source of active transposase, for example, helper plasmid, RNA, transposase protein or integrated transposase source. Transformants containing the desired specific excision product, as shown in Figure 2C, can then be selected.

While efficiency in the second, excision step is less critical, as large numbers of individuals can readily be screened, it is preferred that the excision preferentially removes the short transposon, rather than a long one, as frequently as possible. The present invention provides such a system. piggyBac elements with suitable deletions or mutations in their internal inverted repeat are competent for transposition but not for excision, or have reduced excision rates. Use of such a modified end as the distal end will therefore bias the excision reaction towards utilisation of the proximal end as a higher proportion of the total of excision events.

Thus, it is preferred to use, as a terminal repeat, a repeat having a deletion of no more than 50%, or mutation or inversion that disables no more than 50% of the repeat. It is preferred that such a compromised repeat be in the same orientation as the minimal repeat, where there is only one. These repeats are readily transposed, but are not readily excised after transposition, thereby biasing the excision reaction towards the minimal repeat. Where more than one minimal

repeat is used, in opposing orientations, then more than one compromised repeat may also be used.

The compromised repeat is generally preferred to correspond to a full length repeat, other than in respect of the deletion or mutation.

In relation to either aspect of the present invention, it is possible that subsequent exposure to transposase will simply lead to excision of the entire sequence, in which case the organism is effectively unaltered from the original. The alternative provides the DNA of interest in combination with one full length repeat, but no corresponding inverted repeat, so that the insert is no longer part of a transposable element.

It will also be appreciated that the sequences with which the repeats are associated need not correspond completely to the original sequences found in the naturally occurring transposons, and that variation and sequence degeneracy are encompassed within the scope of the present invention. In particular, it is preferred that any original sequence associated with the repeats of the present invention have at least 70% homology with the corresponding natural sequence, more preferably at least 80%, more preferably at least 90%, and particularly 95% or above, especially 100%.

In order to ensure that the resulting organism has been transformed in the manner desired, it is preferred to utilise appropriate markers. These may be used in any manner suitable to inform the skilled person as to the status of the transformant. For example, markers may be associated with the DNA of interest in order to demonstrate that the organism has successfully been transformed. Markers may be incorporated in the areas between repeats that are to be deleted, so that the initial transformant containing the full length sequence can be detected. Any suitable combination of markers may also be used.

It is generally preferred that markers be selectable, either positive or negative, and suitable examples are illustrated hereinunder.

The use of any transposable element is envisaged, but class II elements, such as Hermes, hobo, Minos, and mariner, are preferred, owing to their relatively high fidelity during transposition, and the piggyBac element, which is known to use the distal element of a repeated

pair at relatively high frequency [Elick, et al., (1997), Mol. Gen. Genet. 255, 605-610], is particularly preferred.

Suitable transposable elements will be known to the skilled person and although reference is made to piggy Bac elements in several examples of the present invention, it will be appreciated that the present invention is not limited thereto and that such reference was made purely for the purpose of exemplification of the invention.

The present invention also provides a method of creating a new marker, preferably by a rearrangement such as that discussed below. Furthermore, the present invention also extends to a method of removing a conditional lethal. If one or both of the flanking transposons contained a conditional lethal gene, one could use this as a negative selection for the original element, and thereby select the desired excision product, even if quite rare. This is equivalent to the use of visible markers, but simpler and requires less effort, so that more transformants can be screened.

The present invention has been shown to work in insects as these are useful and well understood models for genetic transposition. However, it will be appreciated that the present invention will also be useful in a whole range of organisms for transposition into a DNA genome. Accordingly, the present invention is useful in mammals, but also in plants, fungi, and even prokaryotes and viruses.

The present invention, therefore, also provides a method for transforming an organism, comprising exposing replicative cells or tissue of the organism to an element according to the present invention under conditions effective to incorporate the element into the genome thereof and, subsequently or simultaneously therewith, providing conditions suitable to excise a transposon from the genome, and selecting an organism, or cells or tissue therefor, comprising the DNA intended for insertion lacking repeats in at least one orientation. Preferably, the organism is a mammal, a plant, a fungus, a prokaryote, such as bacteria or a virus.

The present invention also encompasses an organism obtained in accordance with the transposable element and method discussed herein. Preferably the organism is an insect.

The final transgenic line preferably has no transposon DNA whatsoever associated with the insertion, unless this has been deliberately incorporated into the DNA of interest. Since there

is no known upper or lower limit for the length of transposition-competent piggyBac elements that would constrain the length of the DNA of interest, the present invention provides a method of very general utility for inserting DNA into genomes of cells or organisms. However, we generally prefer that the total length of the initial piggyBac transposon be in the range 3-25kb, as this corresponds approximately to the range of piggyBac transposons commonly used for germline transformation in insects.

There is no lower limit to the amount of DNA that can be inserted by the overall procedure, after the flanking transposons have been excised. The initial insertion will retain the target site specificity of the original element, such as TTAA for piggyBac, with some apparent preference for (A/T)N(A/T)TTAA(A/T)N(A/T), which may also be written as WNWTTAAWNW (SEQ ID NO. 20) where "W" denotes A or T. Precise excision of the elements will resolve this to a duplication of the TTAA, flanking the DNA of interest, which can be as short as a single nucleotide. In the event that zero nucleotides are inserted, only the TTAA duplication remains. The insertion of larger fragments is generally preferred.

A suitable example of a small insertion is a stop codon. Insertional mutagenesis using transposable elements is a well known method for genetic screens of various types. However, interpreting the phenotype may be complicated by the presence of the transposon, with its associated markers, promoters and other elements. A short insertion, such as TTAA or CTAG, which provides a total sequence between the piggyBac ends of TTAATTAATTAA (SEQ ID NO. 1) and TTAAGTAGTTAA (SEQ ID NO. 2), respectively, allows the insertion to be resolved to a TTAA duplication with this four base insertion. TTAATTAATTAA (SEQ ID NO. 1) and TTAAGTAGTTAA (SEQ ID NO. 2), in these examples, provide stop codons in all three frames in both directions. An insert of zero base pairs provides a frame shift and a stop codon in two frames, although one of these is already present in the original TTAA.

The present invention may be used to provide a coding region, such as for a fluorescent protein or a transactivator protein, such as GAL4, GAL4delta, or tTA. An insertional mutant may then be resolved to a fusion protein. As this needs to be in frame and in the correct orientation to function, it is generally preferable to also provide a suitable IRES (internal ribosome entry site) element and coding region, to allow bicistronic expression, or two such in opposite directions to allow bicistronic expression for insertions of either orientation.

The present invention may also be used to insert an enhancer or promoter or a suitable target site for a site-specific recombinase, either with or without a marker and other sequences. Initial experiments with the non-deleted transposon may serve to establish the presence or absence of specific enhancer effects on a particular insertion site, and the viability and other properties of individuals carrying large insertions at this site. Suitable insertions could then be resolved by stepwise excision to the recombinase target site, either with or without a marker and other sequences. This could then be used as a docking site to enable the insertion of other DNA sequences at this pre-determined and partially pre-characterised position. This arrangement has significant benefits in allowing the insertion of heterologous DNA at a predetermined site. If desired, the site can be protected from nearby enhancers by flanking the recombinase target site with suitable insulator elements, for example scs and scs', or HS4.

The present invention may also be used to replace an existing transposon insertion with specific DNA in such a way that the insertion is substantially or completely stable to further exposure to active transposase. This can be performed by replacing the existing transposon with a composite transposon of the present invention. Methods for performing such a replacement have been described [*c.f.* Johnson-Schlitz, *et al.*, *Mol Cell Biol* **13**, 7006-18 (1993); Cabrera, *et al.*, *Genesis* **34**, 62-5 (2002); Sepp, *et al.*, *Genetics* **151**, 1093-101 (1999); Lankenau, *et al.*, *Mol Cell Biol* **16**, 3535-44 (1996); Gonzy-Treboul, *et al.*, *Genes Dev* **9**, 1137-48 (1995); Heslip, *et al.*, *Genetics* **138**, 1127-35 (1994); Gloor, *et al.*, *Science* **253**, 1110-7 (1991)].

Minimal molecular markers are generally characteristic of one aspect of the invention. It will be apparent to the person skilled in the art that PCR-based or other molecular analysis is capable of distinguishing each of the various possible forms at each stage.

However, with respect to both aspects of the invention, it is generally more convenient to provide at least one visible or selectable marker, and this is preferred. Suitable markers are well known to the person skilled in the art, and include: genes encoding fluorescent proteins, including GFP, DsRed and their mutant derivatives; genes encoding drug or antibiotic resistance, such as neomycin phosphotransferase, or hygromycin resistance; and markers capable of functionally complementing a visible mutant in the host organism, such as mini-*white*⁺ or *rosy*⁺ in *Drosophila*, *white*⁺ in *Ceratitis capitata* or *Drosophila cinnabar* complementing *kw*^w in *Aedes aegypti*. It is particularly preferred that at least one such marker be associated with the section of

DNA containing the DNA of interest, allowing the segment to be tracked through the integration and stepwise excision process.

Where the DNA of interest segment is associated with two repeated ends, then each of the two flanking elements may be marked. Double selection may be used to recognise the initial insertion and then track the stepwise excisions. Since the last step has no high-probability alternatives, individuals with the DNA of interest can be recovered in the final step even though they no longer have any associated visible marker. Thus, the present invention provides a novel method for recovering transgenics with no selectable marker.

Visible markers can generally be scored for or against, in other words are generally suitable for either positive or negative selection. Automated, or semi-automated systems for identifying, or identifying and separating individuals are available, and have the potential for screening very large numbers of individuals, for example in the transformation or excision steps. For the excision step, negative selection for the appropriate transposon is desirable, together with positive selection for the DNA of interest, or an associated marker, as appropriate for the precise scheme selected. Negative selection markers are also known that can be used to select against particular individuals carrying them; such a system can be conveniently used to screen, or to help to screen large numbers of individuals for excision of a specific region. Suitable negative selection markers include inducible or repressible lethals, one half of a biphasic expression system such as GAL4/UAS or tTA/tRE, if the other half can be provided separately, dominant temperature sensitive lethals such as the *Drosophila* DTSs, or synthetic ones such as a suitable toxic element operably linked to a heat-shock or other inducible or repressible promoter.

Multiple visible markers can be provided not only by using mutant derivatives of fluorescent proteins, for example, of GFP and DsRed, which are independently distinguishable based on their spectral properties (*c.f.* Horn *et al.*, 2002, *supra*), but alternatively by expressing the same or similar markers in different spatial or temporal patterns. For example, it is readily possible to distinguish between *Drosophila* Act5C-DsRed, which shows ubiquitous expression, particularly clear in the body of larvae and in the adult eye, and Act88F-DsRed, which shows in indirect flight muscles only, and is, therefore, visible in the thorax of late pupae and adults. These markers are, therefore, separated by both their spatial and their temporal patterns and insects carrying one, the other, or both can readily be distinguished.

Where the present invention relates to transposable elements with two repeated ends, though the initial transformation reaction and the first excision reaction have several potential outcomes, of which not all are desirable, the second excision reaction has only one excision product, and this is the desired one. Accordingly, it is possible not to mark one of the flanking transposons, to select by suitable methods the desired products of the first two reactions, and then to conduct the second excision reaction blind, by exposure to transposase, for example, at high concentration, or for several generations of exposure to a jumpstarter element, then identify the desired reaction product by molecular methods. It is also possible to perform both excision reactions in this way, selecting only for the presence of a marker in the intervening sequence. It is, therefore, possible to perform the entire sequence of reactions using only a marker in this region. This embodiment provides for particularly short flanking transposons, and thus for the maximum size of insert for a given initial composite transposon size.

It will be appreciated that though each step of the insertion sequence is described separately, in practice, exposure to transposase may induce several of these steps to occur within one generation, or without the intermediate being specifically identified. This is generally acceptable and may provide a faster route to the desired final structure, and may be encouraged or stimulated by the use of relatively high concentrations of transposase, or the use of hyperactive transposase or *cis*-acting sequences, if desired. It will be understood that suitable use of molecular and/or selectable markers can facilitate this process.

Transposases are necessary to the function of the present invention, but it is not critical as to how they are provided. They may be provided in any suitable manner, as detailed below, and may be inherent in the cell, provided on plasmids or even provided within the element itself, although this is not preferred. The cell may also be dosed with the enzyme or mRNA encoding the enzyme, or even with a virus expressing the enzyme, for example.

Suitable transposase can be provided in any of several forms: injection or electroporation, for example, of a plasmid or RNA encoding the transposase, or of transposase protein itself. A transposase source may also be integrated into the target genome, to provide a 'jumpstarter' construct, or line. This is a preferred method for the excision steps; the jumpstarter element can be combined with the initial insertion by conventional breeding, or by making the primary transformant in the jumpstarter line. Excision products will then be generated spontaneously, without requiring further injection or electroporation. With suitable markers or molecular

analysis, the desired products can readily be isolated, and separated from the jumpstarter, if required, by conventional breeding.

Although jumpstarters provided within the transposons of the invention are not generally preferred, it is possible to provide a coding sequence therefor in a shorter internal transposon that it is desired to excise. Thus, it is possible to generate a jumpstarter line through insertion of an autonomous element, followed by its resolution by self-catalysed excision.

Transposition is thought to work by a number of mechanisms. One such mechanism involves “scanning” by the transposase. In this mechanism, the transposase binds to one repeat, probably a specific repeat, and scans along the DNA looking for the other repeat. This scanning may or may not be directional. However, where more than one repeat can be recognised by the transposase, as shown for instance in Figure 2, the nearer of two duplicated repeats will be used much more frequently than the other repeat.

This would not appear to be the case for *piggyBac*, which is, therefore, thought not to use such a scanning mechanism, but probably is the case for other class II transposons.

Therefore, in order to enhance the rate of transposition of the full length element comprising the gene of interest, minimal ends may be used, as described above, or an internal rearrangement of the element after insertion can be induced.

The rearrangement or inversion approach starts with a plasmid with only two functional transposable elements, where one transposable element includes the DNA of interest, whereas the other uses the “internal” ends and excludes the DNA of interest, see Figure 3A. After transposition into the genome and recombinase-induced rearrangement/inversion, the initial transposon is converted into a sequence comprising the gene-of-interest and two flanking transposable elements, together with a longer transposable element comprising all three, see Figure 3B. However, under the scanning mechanism, the two shorter flanking transposable elements should then be much more readily mobilised, leaving only the gene-of-interest and one (Figure 3C), but preferably neither (Figure 3D), of the flanking elements remaining.

Inversion is when a chromosomal segment is excised and reinserted in the same place but turned 180 degrees from its original orientation, so that the gene sequence for the segment is reversed with respect to that of the rest of the chromosome.

Suitable systems for such an inversion are well known in the art, such as Cre/lox and Flp/Frt recombinase systems. The cre recombinase is a product of lambda phage in *Escherichia coli*, and the FLP recombinase is an enzyme native to the 2 micron plasmid of *Saccharomyces cerevisiae*. These recombinases alter the arrangement of DNA sequences in very specific ways. The FLP recombinase, for instance, is active at a particular 34 base pair DNA sequence, termed the FRT (FLP recombinase target) sequence. When two of these FRT sites are present, the FLP enzyme creates double-stranded breaks in the DNA strands, exchanges the ends of the first FRT with those of the second target sequence, and then reattaches the exchanged strands. This process leads to inversion or deletion of the DNA which lies between the two sites. Whether there is an inversion or deletion depends on the orientation of the FRT sites: if the sites are in the same direction, the intervening DNA will be deleted, but if the sites are in opposite orientation, the DNA is inverted.

Recombination using wild-type lox or FRT sites is reversible. However, the use of directional recombinase sites, e.g. mutants of lox such as lox66 and lox71, which will recombine with each other in an essentially irreversible reaction, is also envisaged, and generally preferred as these will prevent re-inversion whereby the inversion cassette is subjected to a second round of inversion, such that it is returned to its original orientation. Therefore, the use of directional recombinases biases the inversion to give only the desired product. Other recombinases, whose wild type targets are non-identical and have an essentially irreversible reaction, are also known, e.g. phage C31 integrase acting on attP and attB sites.

Therefore, it is envisaged that the element comprises two external opposed inverted repeats bounding an inversion cassette. The cassette comprises the DNA for insertion into a host genome, together with two inverted repeats and two inversion sites. The DNA for insertion into a host genome is flanked on either side by one of the inverted repeats, each inverted repeat being further flanked by an inversion site, such that reading 5'-3' along one strand of DNA in the cassette, there is provided an inversion site such as FRT, a first inverted repeat, the gene or other DNA of interest, followed by a second inverted repeat and finally by the second inversion site.

The cassette is capable of inversion within the transposed element *in situ*, in the presence of a recombinase. Once the full length transposon has been transposed into the host genome, the recombinase induces inversion of the DNA between the inversion sites.

The result of this is that that following inversion, the two inverted repeats, flanking the DNA of interest, each separately form a new pair of opposing inverted repeats. They do so with one of the external inverted repeats, thus forming a new pair of opposing repeats that are being excisable by a transposase *in situ*, thus leaving the DNA of interest without flanking transposon-derived repeats in the host genome.

The inversion sites, are preferably suitably orientated with respect to each other to allow inversion rather than deletion of the DNA therebetween. For instance, with the Flp/FRT system commonly known to in the art and used here as a non-limiting example only, are preferably aligned in opposite orientations to allow inversion rather than deletion.

The inversion sites are preferably suitably orientated with respect to each other to allow inversion rather than deletion of the DNA therebetween,

As with the transposase the recombinase may, preferably, be encoded by the transposable element of the present invention or, even more preferably, separately therefrom.

Clearly, this recombinase-based inversion approach requires an additional step. The efficiency of this step could be increased by using mutant recombinase target sequences, which make the reaction essentially irreversible. A good marker system would also be useful to detect such rearrangements and is provided in accordance with the invention. Here, for instance, the recombinase target sites are embedded in two different markers, such that following rearrangement, two new markers are created.

For example, marker 1 might be EGFP under the control of an eye-specific promoter; marker 2 might be DsRed under the control of a ubiquitous or body-specific promoter. The recombinase target site is included in each marker between the promoter and the coding region for the fluorescent protein. The initial transgenic then exhibits green fluorescent eyes and red fluorescent body, while individuals carrying the desired recombination product (which is an inversion of an internal section of the composite transposon) show red fluorescent eyes and a

green fluorescent body. Individuals carrying the desired inversion should therefore be readily detected, even if rare.

Accordingly, the present invention also provides a method of creating a new marker by such a rearrangement, such as the red eyes produced in the above example. Furthermore, the present invention also extends to a method of removing a conditional lethal.

Without being bound by theory, it is worth noting that the 31 base pair spacer found in the right hand (3') piggyBac repeat may wrap around a protein, in contrast to the 3 or 4 base pair spacer in the 5' repeat, perhaps suggesting that the transposon is an evolutionary ancestor of the VDJ antibody variation domain of the MHC, where a similar mechanism is also seen.

Experiment 1: Use Of Minimal Ends

We made construct pLA1025 (see Figure 5 and SEQ ID NO. 3), which includes minimal piggyBac ends (35bp and 63bp bases for 5' and 3' respectively). We generated transgenic *Drosophila melanogaster* carrying this construct by co-injection with a helper plasmid into a white-eyed strain (Handler and James 2000; Handler 2002, below). Potential transgenics were screened for fluorescence characteristic of EGFP and DsRed. pLA1025 contains 4 potential transposons, marked with [Actin5C-EGFP, Actin5C-DsRed2 and Actin88F-DsRed2], [Actin5C-EGFP and Actin88F-DsRed2 but not Actin5C-DsRed2], [Actin5C-EGFP only] and [Actin88F-DsRed2 only], respectively.

If all transposon ends were utilised with equal efficiency, one might expect these 4 transposons to be recovered with equal frequency, or possibly a higher frequency for the shorter transposons which contain only one marker. Of 5 independent transgenic lines recovered from 85 G0 adult injection survivors, all showed all three markers, and therefore correspond to the desired composite transposon. This indicates that the utilisation of ends and nature of the recovered transposons can be biased in a desirable way by using internal ends that are less readily utilised for plasmid-to-germline transposition.

A diagrammatic representation of the pLA1025 construct is shown in Figure 5 and its predicted sequence given in SEQ ID NO. 3.

Experiment 2: Resolution Of Four-End Constructs To Give Transgenic Strains In Which DNA Of Interest Is No Longer Flanked By Transposon DNA.

We made construct pLA1125 which includes 4 piggyBac ends, two each of 5' and 3' ends. pLA1125 is shown diagrammatically in figure 6 and its predicted sequence is shown in (SEQ ID NO. 4)

We generated transgenic *Drosophila melanogaster* carrying this construct by co-injection with a helper plasmid into a white-eyed strain (Handler and James 2000; Handler 2002, below). Potential transgenics were screened for fluorescence characteristic of DsRed. We recovered transgenics in the progeny of 12 out of 227 adult G0 injection survivors. PCR analysis for the presence of each of regions 1, 2 and 3, and their contiguous presence, showed that all these transgenic lines contained a composite transposon of the expected structure.

This demonstrated that, although pLA1125 contains 4 potential transposons, transgenics containing the desired composite transposon (PB5'-ZsGreen-PB3'-DsRed2-PB5'-AmCyan-PB3') could be recovered at a useful rate. We found that neither ZsGreen nor AmCyan could be reliably scored by fluorescence microscopy in these transgenic flies, despite the presence of the coding regions for these proteins as analysed by PCR, so subsequent experiments used fluorescence microscopy as a reliable determinant of the presence of DsRed only. Parallel experiments with other constructs indicated that the *OpIE2* promoter fragment used to drive expression of ZsGreen and AmCyan in LA1125 gives only weak expression in *Drosophila*, furthermore the AmCyan region of LA1125 contains a frameshift likely to reduce or eliminate production of functional AmCyan protein.

One of these transgenic LA1125 lines (line12) was selected for further study. We determined the insertion site by obtaining flanking DNA and sequencing, then comparing this sequence with the known genomic DNA sequence of *Drosophila melanogaster* from the *Drosophila* genome project. This showed that the composite LA1125 element had inserted into the sequence located on chromosome 2. This is shown in Figure 7 where the site of insertion is written in capital letters and underlined, and in (SEQ ID NO. 5).

We exposed the integrated LA1125 line 12 (henceforth "1125-12") to piggyBac transposase by crossing to piggyBac "jumpstarter" lines, either pCasper-hs-orf (see <http://piggybac.bio.nd.edu/>), or pHer{3xP3-ECFP, hsp70-piggyBac} (Horn, N et al. 2003). Progeny of these crosses were heatshocked during embryonic and larval development (37°C for 1 hr, 2x per week, starting 0-7 days after egg deposition, at which point parents were removed, stopping when first pupae were observed in vial). Double heterozygous (1125-12/+ and jumpstarter/+) F1 adults were selected, based on marker expression.

Somatic transposition

Extracts from pools of 5 double heterozygous (1125-12/+ and jumpstarter/+) adults, or from individual flies, were analysed for excision of the flanking simple transposons by PCR, see Tables 1 and 2 below. Precise excision of either element would bring the flanking genomic DNA closer to the central DsRed gene. This would lead to a characteristic, predictable, reduction in size of a PCR product using Dro-12m-4 + su-a5c-r1 primers or Dro-12m-1 + Diag-droso primers for PCR, respectively detecting excision of the ZsGreen or AmCyan simple transposon.

Primer's name	Predicted size of PCR product from 1125-12	Predicted size of PCR product from 1125-12 following deletion of ZsGreen simple transposon
Dro-12m-4 + su-a5c-r1	3058 bp	277 bp

Table 1

Primers	Predicted size of PCR product from 1125-12	Predicted size of PCR product from 1125-12 following deletion of AmCyan simple transposon
Dro-12m-1 + Diag-droso	2913 bp	268 bp

Table 2

In all pools and all individuals analysed, we observed the characteristic bands expected for precise excision of each simple transposon (Figure 8). We did not observe such bands from similar extracts of flies from the 1125-12 line which had not been exposed to transposase. We therefore conclude that it is possible to eliminate each of the two flanking transposons from the composite 1125 transposon, while retaining the central region. We further conclude that this excision is substantially dependent on the presence of piggyBac transposase, as expected. Since the PCR band is of the expected size for a precise excision, and piggyBac excisions are known to be precise in the large majority of cases, we conclude that precise somatic excision of the flanking elements is occurring at a readily detectable rate.

Figure 8 shows the PCR detection of somatic excision of the flanking transposons in F1 adults flies after exposure to transposase, wherein:

M = DNA marker (SmartLadder, Eurogentec)

A = PCR for the excision of ZsGreen simple transposon

B = PCR for the excision of AmCyan simple transposon

A1 and B1 = template DNA from a pool of 5 F1 flies which were exposed to transposase

A2 and B2 = template DNA from a single F1 fly which was exposed to transposase

A3 and B3 = template DNA from a single fly which had no exposure to transposase

Germ-line transposition

In order to determine whether the flanking transposons could be eliminated in the germ-line, to generate gametes containing the excision products and hence a heritable excision product, double heterozygous (1125-12/+ and jumpstarter/+) F1 adults were crossed to white-eyed (w^{1118}) flies. Pools of 7 F2 progeny flies, selected for the presence of the DsRed marker by fluorescence microscopy, were obtained. Extracts from these pools were analysed for the presence of the specific excision products as above. Several pools yielded specific PCR bands as for the somatic excision assay above. One example is shown in Figure 9. This indicates that germline excision can occur as for somatic excision above.

Figure 9 shows the PCR detection of germline excision of ZsGreen simple transposon, wherein:

M = DNA marker (SmartLadder, Eurogentec)

1 – 7 = DNA extracted from different pools of 7 F2 1125-12 flies (with DsRed)

Dro-12m-4 and su-a5c-r1 primers were used in these PCR reactions

192 individual flies were analysed for excision by using the primers which had been used to detect the somatic and the germline excision of ZsGreen and Amcyan simple transposons. These flies were also analysed for the presence of ZsGreen (su-zsgreen-f + su-zsgreen-r) and Amcyan (su-amcyan-f + su-amcyan-r) simple transposons and for connections between the central DsRed region with ZsGreen (Diag-rab-b-g + su-a5c-r1) and with AmCyan (Opie2-nhe-R + Diag-droso) simple transposons.

The somatic excision data, and the data from pools of F2 individuals, clearly indicates that precise excision of the flanking transposons can occur. However, in a careful PCR-based analysis of four of the individual F2 flies showing excision events, we were unable to amplify a

diagnostic flanking DNA band of the type used to show precise excision in somatic cells. However, these flies had clearly lost one or both of their flanking transposons, as judged by the inability to amplify unique regions from one or both of these transposons (primers su-zsgreen-f and su-zsgreen-r for the ZsGreen transposon and su-amcyan-f and su-amcyan-r for the AmCyan transposon), while being readily able to amplify a unique region from the central DsRed region (primers su-red-f and su-red-r) and indeed to observe fluorescence characteristic of DsRed in these flies.

We conclude that the excision event that removed the flanking simple transposons was associated with a rearrangement, possibly a deletion of flanking genomic DNA but, more likely, a transposition of the composite transposon such that it is no longer associated with the original flanking DNA but is now inserted at a new site in the genome. The ability to recover remobilised, products of this type is an advantage, as it indicates that an initial primary transformant can be simultaneously relocated to another site and stabilised by elimination of one or both flanking transposons, which saves a considerable amount of time over the multi-generational breeding scheme that would be required to obtain such a product if each step had to be performed sequentially.

We further analysed the two DsRed-positive flies in which neither ZsGreen nor AmCyan appeared to be present. PCR using primers PB5-sh-srf-asc-1 and PB5-sh-not-hind-2, which amplify a section of piggyBac, showed that no such piggyBac DNA is present anywhere in these flies (Figure 10). The ZsGreen and AmCyan transposons have not, therefore, rearranged or transposed to another part of the genomes of these flies, rather they have been completely lost, presumably by excision in the germ-line of the F1 parent. We have therefore demonstrated that it is possible by this method to generate transgenic flies in which the inserted DNA is not associated with the ends of a transposon.

Figure 10 shows the results of PCR analysis of piggyBac sequence in DsRed stable transformants, wherein:

M = DNA marker (SmartLadder, Eurogentec)

1 = DNA from LA1125-12 flies unexposed to transposase used as positive control

2 = DNA from fly 70 (has neither ZsGreen nor AmCyan)

3 = DNA from fly 200 (has neither ZsGreen nor AmCyan)

Markers

The ZsGreen and AmCyan open reading frames were used only as molecular markers in these experiments. Use of visible markers would reduce the amount of PCR required, as phenotypic scoring for the presence of such markers, e.g. for the production of fluorescent proteins, could be used to infer excision and other events. However, we have clearly demonstrated that this is not necessary, that these events occur at sufficient frequency that molecular-based screening alone is sufficient to recover the desired types, e.g. excision products. Two variants of LA1125 are therefore proposed based on the above data: (i) an equivalent construct with visible markers in the flanking transposons and (ii) an equivalent construct with shorter flanking transposons containing no such visible markers. The second of these options should give higher transposition and excision frequencies, as shorter transposons are generally thought to be more active. A short segment of unique sequence in each of the flanking transposons would be helpful for the molecular analysis. Such a construct is also the product of a Flp/Frt, Cre/lox or other similar inversion system following recombinase-mediated inversion of the central section of the construct described.

Vector

pLA1125 provides several unique sequences into which additional DNA could be inserted. Particularly convenient in this regard are the *AscI* and *SrfI* sites in the central section. pLA1125, and the variants described above, therefore provide vectors of general utility for the method of the invention.

Primers

Primers are given below in Table 3

Primer Name	Sequence
Dro-12m-1	gccagtcctgcagcttcagcgtgtg (SEQ ID NO. 6)
Dro-12m-4	aaaggtatgatggtgaagtatccgc (SEQ ID NO. 7)
Diag-droso	ccttctttgttgaatccagatcctgc (SEQ ID NO. 8)
Diag-Ra-b-g	gctcctgggcaacgtgctggttg (SEQ ID NO. 9)
Opie2-nhe-R	ggtggctagcttgcgcttcttcttggtggg (SEQ ID NO. 10)

PB5-sh-srf-asc-1	ggtggcgcgcccaattgccgggcttttaaccctagaaagatagtctgcgt (SEQ ID NO. 11)
PB5-sh-not-hind-2	gcccaagcttgcgccgcgtcatttgactcacgcggctcg (SEQ ID NO. 12)
su-a5c-r1	gattgattcagagttgatgccattcatg (SEQ ID NO. 13)
su-amcyan-f	ccagacctccacctcaaggtgacc (SEQ ID NO. 14)
su-amcyan-r	ttgtaggaggtgtggaactggcatctg (SEQ ID NO. 15)
su-red-f	caacaccgtgaagctgaaggtgacc (SEQ ID NO. 16)
su-red-r	cttgccatgtagatggacttgaactcc (SEQ ID NO. 17)
su-zsgreen-f	caagcaggccatcaacctgtgc (SEQ ID NO. 18)
su-zsgreen-r	gacttggcctgtacacggtgtcg (SEQ ID NO. 19)

Table 3

References for Experiments 1 and 2

Handler, A. (2002). "Use of the *piggyBac* transposon for germ-line transformation of insects." *Insect Biochem. Mol. Biol.* **32**: 1211-20.

Handler, A. and A. James (2000). *Insect transgenesis: methods and applications*. Boca Raton, CRC Press.

Horn, C., O. N, et al. (2003). "piggyBac-based insertional mutagenesis and enhancer detection as a tool for functional insect genomics." *Genetics* **162**(2): 647-661.

Claims:

1. 1. A transposable element comprising at least four inverted repeats, forming at least two pairs of opposing pairs of inverted repeats, the element comprising DNA for insertion into a host genome, the DNA being located between two pairs of opposing repeats such that excision by a transposase or transposases of said pairs, *in situ*, is effective to be able to leave said DNA integrated into the host genome, without the presence of flanking transposon-derived repeats derived from said transposable element.
2. A transposable element according to claim 1, wherein the DNA for insertion into a host genome is a gene for expression in the host.
3. A transposable element according to claim 1, wherein the DNA for insertion into a host genome is a promoter or enhancer sequence.
4. A transposable element according to claim 1, wherein the DNA for insertion into a host genome is a stop codon or is sufficient to bring about an in frame stop codon.
5. A transposable element according to any preceding claim, wherein the inverted repeats are piggyBac repeats.
6. A transposable element according to any preceding claim, having four inverted repeats.
7. A transposable element according to any preceding claim, wherein the inverted repeats are homologous.
8. A transposable element according to any of claims 1-6, wherein pairs of homologous inverted repeats are heterologous to other pairs of inverted repeats.
9. A transposable element according to any preceding claim, wherein one or more of the inverted repeats is a minimal non-terminal repeat.
10. A transposable element according to any preceding claim, comprising at least one genetic marker.

11. A transposable element according to any preceding claim, wherein the element comprises two external opposed inverted repeats, one on each side of an inversion cassette, the cassette comprising;

the DNA for insertion into a host genome, two inverted cassette repeats and two inversion sites, the DNA for insertion into a host genome being flanked on either side by one of the inverted cassette repeats, each inverted cassette repeat being further flanked by an inversion site;

the cassette being capable of inversion within the transposed element *in situ* in the presence of a recombinase, such that following inversion, the two inverted cassette repeats flanking the DNA for insertion into a host genome each separately form a further pair of opposing inverted repeats with one of the external inverted repeats, the further pairs of opposing repeats being excisable by a transposase *in situ* to leave said DNA without flanking transposon-derived repeats in the host genome.

12. A transposable element according to claim 11, wherein the inversion sites are recognised by inversion-inducing recombinase.

13. A transposable element according to claim 12, wherein the inversion sites are recognised by the Flp/FRT or Cre/lox inversion systems.

14. A transposable element comprising at least three inverted repeats, at least one of which is inverted in relation to the others, wherein at least one non-terminal repeat is a minimal repeat.

15. A transposable element according to claim 14, comprising DNA for insertion into a host genome located between the minimal repeat and a repeat having the same orientation as the minimal repeat.

16. A transposable element according to claim 14 or 17, wherein the DNA for insertion into a host genome is preferably flanked by two pairs of opposing repeats excisable by a transposase *in situ* to leave said DNA without flanking repeats in the host genome.

17. A transposable element according to claim 16, wherein each of the repeats bounding the DNA for insertion into a host genome is a minimal repeat.

18. A transposable element according to any of claims 14-17, wherein at least one repeat distal to the DNA for insertion into a host genome in relation to a minimal repeat in the same orientation has an internal deletion or is otherwise compromised over up to 50% of its length.
19. A transposable element according to any of claims 10 and 14-18, comprising at least one genetic marker associated with an identifiable step in the transposition/excision process.
20. A transposable element according to claim 19, wherein the marker is associated with the DNA for insertion into a host genome.
21. A transposable element according to any of claims 14-19, comprising as a terminal repeat, a repeat having a deletion of no more than 50%, or mutation or inversion that disables no more than 50% of the repeat.
22. A transposable element according to any preceding claim, wherein the element is a class II transposable element.
23. A transposable element according to any preceding claim, wherein the transposase is encoded within the transposon.
24. A method for transforming an organism, comprising exposing replicative tissue of the organism to an element according to any preceding claim under conditions effective to incorporate the element into the genome thereof and, subsequently or simultaneously therewith, providing conditions suitable to excise a transposon from the genome, and selecting an organism, or tissue therefor, comprising the DNA intended for insertion lacking repeats in at least one orientation.
25. A method according to claim 24, wherein the transformant organism is exposed to a source of active transposase.
26. A method according to claim 25, wherein the source of active transposase comprises a helper plasmid or RNA encoding the transposase, or a transposase protein or integrated transposase source.

27. A transformant organism obtained in accordance with any of claims 24-26.
28. A transformant organism according to claim 27, wherein the organism is an insect.
29. A transposable element according to any of claims 10 and 19-20, wherein the marker is a conditional lethal.
30. A transposable element according to any claim 12, wherein the inversion sites are recognised by a directional recombinase, the recombinase-mediated inversion being essentially irreversible.
31. A transposable element according to any claim 30, wherein the inversion site is lox66 or lox71.

1 / 10





Figure 1

2 / 10

**Figure 2**

3 / 10




Figure 3

4 / 10




Figure 4




Figure 5

6 / 10




Figure 6

7 / 10

167161 gaaatcgattgcggaaaccattgcttagtTTTTgattgacctttgtttgaaaacaaaat
167221 atatgtataaaaggaattatttgaaagtacgagcggtttgttgatcatattggtgggattt
167281 ttggctatttcattttaattaagtTTTTgaatttaaagtgttcatttagctacattca
167341 gaaatattgttttgaatttaaaaaatttacagtgtgaatgtcttgaattttcaattgg
167401 aaaaaaaggatgatggttgaagtatccgctttgtttattacagtagactggtcacact
167461 gccagctgctaaataatagaaaacagtctgcgtggacaaacatttaaTTAAaatgaatg
167521 taagcactttttaacgaaatctttgggaatatctcgctcatcagcattttatttgagca
167581 ggagtccgagtcccatcgacatacacacgctgaagctgcaggactggctgatcagccg
167641 cagaattgtgccagaatgtgcagcaggagcttagggagatccacaggaagattagcaa

Figure 7






Figure 8

9 / 10

**Figure 9**

10 / 10

**Figure 10**

SEQUENCE LISTING

<110> Oxitec Limited
 <120> STABLE INTEGRANDS
 <130> WPP88351
 <150> UK 0315379.8
 <151> 2003-07-01
 <160> 20
 <170> PatentIn version 3.2
 <210> 1
 <211> 12
 <212> DNA
 <213> artificial
 <220>
 <223> Transposon Inverted Repeat
 <400> 1
 ttaattaatt aa 12
 <210> 2
 <211> 12
 <212> DNA
 <213> artificial
 <220>
 <223> Transposon Inverted Repeat
 <400> 2
 ttaactagtt aa 12
 <210> 3
 <211> 16157
 <212> DNA
 <213> artificial
 <220>
 <223> predicted sequence of pLA1025 construct
 <400> 3
 gggcaattcg ggcgcgctc ttaattaagg cccccctcg agaccatggt cgatcgacaa 60
 acacgaaaat acattcctgg acagacctag gggtttgctc ttttttcttt gcaggatctg 120
 gattcaacaa agaaggaaaa atctacggag ttggaacct catttattgc acgaagaaac 180
 ttagagagaa agactgatgc ctaataaaga ctcaagctga ttgcaaaaac aatataaagt 240
 tgtaccttat tgcggtgtca tcgtaaaacc aagatgctta aaagggatta ggcccactta 300
 aattactgcg aaaaatttta ttacaaatt tctcacatga cttccagata aactaattaa 360
 ctttgtgtgt caagggtgct tttccggcag aaaagtcaat ttgcaagtga tttagggtttt 420
 tttaatgact atctttgcaa ttgaattggg tactattaga tattgaatat accaatattg 480
 aggtataata caaatacaac aacatttgac ctcggaact atcagaagta catgtaggtc 540
 gggaacatta gggaattctg gcgataagt tactaattat gctaagctat cgcagaacct 600

cattaagact	tgagctgcag	tgttttgcag	gtacttatca	tatttacagt	attaatgtat	660
gcagtattat	attaattcta	aataagaaag	tcaggagaag	gatattcaaa	actgaagaat	720
taatgtacat	tagttttggt	tatttagattt	tcaatattgc	agcatagaat	atgtgtaagt	780
agtggagagc	taaacgcgct	tttcagaacg	ttcatgctaa	ttgctcatgg	atctcacagg	840
aacagggtgg	ggcggccctc	ggtgcgctcg	tactgctcca	cgatgggtga	gtcctcgttg	900
tgggaggtga	tgccagctt	ggcgctccacg	tagtagtagc	cgggcagctg	cacgggcttc	960
ttggccatgt	agatggactt	gaactccacc	aggtagtggc	cgccgtcctt	cagcttcagg	1020
gccttggtgg	tctcgccctt	cagcacgccg	tcgcgggggt	acaggcgctc	ggtggaggcc	1080
tcccagccca	tggtcttctt	ctgcatcacg	gggcccgcgg	aggggaagtt	cacgccgatg	1140
aacttcacct	tgtagatgaa	gcagccgtcc	tcaggggagg	agtcctgggt	cacggtcgcc	1200
acgccgccgt	cctcgaagtt	catcacgcgc	tcaccattga	agccctcggg	gaaggacagc	1260
ttcttgtagt	cggggatgtc	ggcggggtgc	ttcacgtaca	ccttgaggcc	gtactggaac	1320
tggggggaca	ggatgtccca	ggcgaagggc	agggggccgc	ccttggtcac	cttcagcttc	1380
acgggtgtgt	ggccctcgta	ggggcgggcc	tcgccctcgc	cctcgatctc	gaactcgtgg	1440
ccgttcacgg	tgccctccat	gcgcaccttg	aagcgcatga	actcgggtgat	gacgttctcg	1500
gaggaggcca	ttctagccat	tttggttcta	gatccggggt	ctctggatta	gacgactgct	1560
ggctgatgga	gcggctttgt	gtcgggagga	gtatccacac	agcacaagaa	ctcaaacggt	1620
agtgatatga	aaactggtcc	cgcagcccg	tttataaagc	ccgcttagcg	ctaaaacctt	1680
atcagccgca	atgctctccg	ctctcaagtc	gcgttcaaaa	cttttaccat	atatgggtat	1740
gaattgttcg	ccgattgggt	tttcatttac	ggagaatttc	ctccgcaact	gggtgttcgt	1800
gcttgtgtgt	gtcttgata	tcgccatgaa	gagtgtactt	ccatgggtcc	tatcagttgc	1860
ggcatgtgat	tcaatttggg	ggttactcat	attcgatact	tcagctgggt	atctttatgc	1920
tgtgatggtc	acgggtgctt	tacacatttt	ttccccaa	actcattgta	tgccaaaaca	1980
tacatatgta	catgtgttta	tagtgggttt	attagtaatg	tccagcgata	ttaaaacatt	2040
acagttaa	aatattcata	ttagcttacg	tttactgcta	tctgcacttc	aaattgggtta	2100
ttcagggaca	tttgtgcaac	acacatcttc	tagtaggggc	atcactagtc	gtagaaaaca	2160
cacctaa	aagccggttt	taacaataac	ggcgaaacag	ctgccagacg	agagaaaagt	2220
acacagcaag	aagcatttct	ataatgatgc	cgctcctaa	atgcagcatt	tttgctcaaa	2280
ttgatcat	cgttgcatcg	ctcagggtgt	atagtatttg	gctattgaat	gtgattgcta	2340
tcttaatagt	tttgaatagt	aatagtgcaa	atattcattc	ataaaaaata	ggtaatcgat	2400
ttctcagtg	cagtctatgc	gtttgcgcaa	acaggagacg	aagagagaga	cgcctcacac	2460
acaagcacat	gcgtactacg	ctggcctact	ccgtactctt	atgtgtcgtc	ctctgcacgc	2520
tctgcttggc	ggcagttctt	ccagaggctg	tgacgtcagg	taagctat	cgtctcggtg	2580
tgctgttagc	ccgtaacgat	aagctgtccg	attcctaatt	ctctttcaaa	taatttggtt	2640

tattttttttt tttgtgtgtt ttgaagtcaa acagcttcat aaacaatatt aaatatcgca	2700
attgttattg caaatgcatt cgtatcaggc gccttgatgt ttttcggcca aaacttcgtc	2760
tgctctttgt cagcgtaatt gcgtttgccc gtcgtgccgt cttgctcgca cgcgcttgca	2820
aacaaaaagt gaaagagggg aagccactgt gttgttgctt ttgttgctat cgttggtgc	2880
acacacatgg caataagtaa ccatattgta gacttattta ttttagcggt ttcgttgact	2940
aatctttgtt taaataattg cctatttcga gccaaaatga cgcataaagg acacggcaac	3000
atgcggtgcg ttgagcatgc gtacgagatg gcgctagaga gagcgagata aaaagaaaaa	3060
aacaaccggc gagtggcgag acacgaataa tacacacacg cacacatggg caaaggagtg	3120
gctccaacgt cgggtgacgtc gcgcccagcc acagaccccg ccacgccccg cccaccctt	3180
ccttcgctg ccccgcaccc ctttccttcc actaccgttt tgccgcattt ggtattgcc	3240
tctcgctttt ttggctttac ttgcttttt gaataagcca agaaactgaa atcaaagcgg	3300
acagtccgaa aattgaaact attccaaaaa acaacaaccg cacatgcact gcgttcggca	3360
aaagacgcag aaatcagcct gtggtctggc atcatcttaa aatatcatca gtatcatcat	3420
tgccattata aatacataag aaaatgagaa tacattaaag gtatatacat acaatctttg	3480
tcatagtcaa catattttgc aacaaaatta tctatagata acatgttttt gtagaaaatt	3540
ccactaactg attttcgtat aaatacaaat tattctcatt cttcaaaata ctcgccgacg	3600
tcatcgcgaa attcctcagc attcttcgac aaatgaagag agacagcgag tgacacaatc	3660
ttatcttacc gacgagtctt tgaaaatggc cccgggaacg gtgaaaaaca tatagccaag	3720
aatggagcga tcgccaggcg atggaaattg gcttcaaatt gctccgaata tggccagttg	3780
gtcagttggc attggattgt ggagagatac tagatcagct cgttctcctt ctcctccgca	3840
ttctcctact cgcgcgaaga cgcttctttg atcgcttctt tctcgaatgt acatattgat	3900
atgtatttgg tgtttggcga cgatcagcca gctttaatgg ccataaaacg tgtcttggtt	3960
tgtggggccac ataacaggca gcgagatctg gaaaaatgat gtgacagtgg aaatgagaag	4020
taggtgcata tgcaaagatt tgattcagag ttgatgccat tcatgatatt ttttagaagt	4080
atcatttgtg tttttagaat atagaattct agtatgtatg taagttaata aaaccattt	4140
ttgcggaaaag tagataaaaa aaacattttt tttttttact gcactggata tctgcggccg	4200
gccagtttaa ccctagaaaag ataatcatat tgtgacgtac gttaaagata atcatgcgta	4260
aaattgacgc atgggtacca gatatccagt gcagtaaaaa aaaaaaatgt tttttttatc	4320
tactttccgc aaaaatgggt ttattaact tacatacata ctagaattct aaaaaaatc	4380
atgaatggca tcaactctga atcaaatctt tgcagatgca cctacttctc atttccactg	4440
tcacatcatt tttccagatc tcgctgcctg ttatgtggcc cacaaccaa gacacgtttt	4500
atggccatta aagctggctg atcgctcgcca aacaccaa atacatcaat atgtacattc	4560
gagaaagaag cgatcaaaga agcgtcttcg ggcgagtagg agaatgcgga ggagaaggag	4620
aacgagctga tctagtatct ctccacaatc caatgccaac tgaccaactg gccatattcg	4680

gagcaatttg aagccaattt ccatcgctg gcgatcgctc cattcttggc tatatgtttt 4740
tcaccgttcc cggggccatt ttcaaagact cgtcggtaag ataagattgt gtcactcgct 4800
gtctctcttc atttgcgaa gaatgctgag gaatttcgcy atgacgtcgg cgagtatttt 4860
gaagaatgag aataatttgt atttatacga aaatcagtta gtggaatttt ctacaaaaac 4920
atgttatcta tagataattt tgttgcaaaa tatgttgact atgacaaaga ttgtatgtat 4980
atacctttta tgtattctca ttttcttatg tatttataat ggcaatgatg atactgatga 5040
tattttaaga tgatgccaga ccacaggctg atttctgctt cttttgccga acgcagtgc 5100
tgtgcggttg ttgttttttg gaatagtttc aattttcggg ctgtccgctt tgatttcagt 5160
ttcttggtt attcaaaaag caaagtaaag ccaaaaaagc gagatggcaa taccaaatgc 5220
ggcaaaacgg tagtggaagg aaaggggtgc ggggcagcgg aaggaagggt ggggcggggc 5280
gtggcgggggt ctgtggctgg ggcgcagctc accgacgttg gagccactcc tttgacctg 5340
tgtgcgtgtg tgtattatct gtgtctcgcc actcgccggt tgtttttttt tttttatctc 5400
gctctctcta gcgccatctc gtacgcagtc tcaacgcacc gcatgttgcc gtgtccttta 5460
tgcgtcattt tggtctgaaa taggcaatta tttaaacaaa gattagtcaa cgaaaacgct 5520
aaaataaata agtctacaat atggttactt attgccatgt gtgtgcagcc aacgatagca 5580
acaaaagcaa caacacagtg gctttccctc tttcactttt tgtttgcaag cgcgtgcgag 5640
caagacggca cgaccggcaa acgcaattac gctgacaaag agcagacgaa gttttggccg 5700
aaaaacatca aggcgcctga tacgaatgca tttgcaataa caattgcgat atttaattt 5760
gtttatgaag ctgtttgact tcaaaacaca caaaaaaaa aataaaacaa attatttgaa 5820
agagaattag gaatcggaca gcttatcgtt acgggctaac agcacaccga gacgaaatag 5880
cttacctgac gtcacagcct ctggaagaac tgccgccaag cagacgatgc agaggacgac 5940
acatagagta gcggagtagg ccagcgtagt acgcagtgtc ttgtgtgtga ggctctctc 6000
tcttcgtctc ctgtttgctc aaacgcatag actgcactga gaaaatcgat tacctatttt 6060
ttatgaatga atatttgac tattactatt caaaactatt aagatagcaa tcacattcaa 6120
tagccaaata ctataccacc tgagcgatgc aacgaaatga tcaatttgag caaaaatgct 6180
gcatatttag gacggcatca ttatagaaat gcttcttgct gtgtactttt ctctcgtctg 6240
gcagctgttt cgccgttatt gttaaaaccg gcttaagtta ggtgtgtttt ctacgactag 6300
tgatgcccct actagaagat gtgtgttgca caaatgtccc tgaataacca atttgaagtg 6360
cagatagcag taaacgtaag ctaatatgaa tattatttaa ctgtaatgtt ttaatatcgc 6420
tggacattac taataaacc actataaaca catgtacata tgtatgtttt ggcatacaat 6480
gagtagttgg ggaaaaatg tgtaaaagca ccgtgaccat cacagcataa agataaccag 6540
ctgaagtatc gaatatgagt aacccccaaa ttgaatcaca tgccgcaact gataggaccc 6600
atggaagtac actcttcatg gcgatataca agacacacac aagcacgaac acccagttgc 6660
ggaggaaatt ctccgtaaat gaaaacccaa tcggcgaca attcataccc atatatggta 6720

aaagttttga acgcgacttg agagcggaga gcattgcggc tgataagggt ttagcgctaa 6780
 gcgggcttta taaaacgggc tgcgggacca gttttcatat cactaccgtt tgagttcttg 6840
 tgctgtgtgg atactcctcc cgacacaaaag ccgctccatc agccagcagt cgtctaattcc 6900
 agagaccccc gatctagaac caaaatggct agcaaaggag aagaactttt cactggagtt 6960
 gtcccaattc ttgttgaatt agatgggtgat gttaatgggc acaaattttc tgtcagtgga 7020
 gaggggtgaag gtgatgcaac atacggaaaa cttaccctta aatttatttg cactactgga 7080
 aaactacctg ttccatggcc aacacttgct actactttca cttatgggtgt tcaatgcttt 7140
 tcaagatacc cagatcatat gaaacagcat gactttttca agagtgccat gcccgagggt 7200
 tatgtacagg aaagaactat atttttcaaa gatgacggga actacaagac acgtgctgaa 7260
 gtcaagtttg aaggtgatac ctttgtaaat agaatcgagt taaaagggtat tgattttaaa 7320
 gaagatggaa acattcttgg acacaaattg gaatacaact ataactcaca caatgtatac 7380
 atcatggcag acaacaaaaa gaatggaatc aaagttaact tcaaaattag acacaacatt 7440
 gaagatggaa gcgttcaact agcagaccat tatcaacaaa atactccaat tggcgatggc 7500
 cctgtccttt taccagacaa ccattacctg tccacacaat ctgccctttc gaaagatccc 7560
 aacgaaaaga gagaccacat ggtccttctt gagtttgtaa cagctgctgg gattacacat 7620
 ggcattggatg aactatacaa ataaggatcc atgagcaatt agcatgaacg ttctgaaaag 7680
 cgcgtttagc tctccactac ttacacatat tctatgctgc aatattgaaa atctaataaa 7740
 caaaactaat gtacattaat tcttcagttt tgaatatact tctcctgact ttcttattta 7800
 gaattaatat aatactgcat acattaatac tgtaaataatg ataagtacct gcaaaacact 7860
 gcagctcaag tcttaatgag gttctgcgat agcttagcat aattagtaac ttatcgcgca 7920
 gaattcccta atgttccga cctacatgta cttctgatag ttgccgaggt caaatgtgtg 7980
 tgtatttgta ttatacctca atattggtat attcaatata taatagtacc caattcaatt 8040
 gcaaagatag tcattaaaaa aacctaaatc acttgcaaat tgactttttc gccggaaaag 8100
 caaccttgac acacaaagtt aattagttta tctggaagtc atgtgagaaa tttgtaaata 8160
 aaatttttcg cagtaattta agtgggccta atccctttta agcatcttgg ttttacgatg 8220
 acaccgcaat aaggtacaac tttatattgt ttttgcaatc agcttgagtc tttattaggc 8280
 atcagtcttt ctctctaagt ttcttcgtgc aataaatgag gttccaaact ccgtagattt 8340
 ttcttctttt gttgaatcca gatcctgcaa agaaaaaaga gcaaaccctt aggtctgtcc 8400
 aggaatgtat tttcgtgttt gtcgatcgac catggtcacc ggtagatata tataacaaga 8460
 aaatatatat ataataagtt atcacgtaag tagaacatga aataacaata taattatcgt 8520
 atgagttaaa tcttaaaagt cacgtaaaag ataatcatgc gtcattttga ctcacgcggt 8580
 cgttatagtt caaaatcagt gacacttacc gcattgacaa gcacgcctca cgggagctcc 8640
 aagcggcgac tgagatgtcc taaatgcaca gcgacggatt cgcgctatct agaaagagag 8700
 agcaatatct caagaatgca tgcgtcaatt ttacgcagac tatcttttcta gggttaagtc 8760

gacatccaca ttgtggccaa gctgtgtgac gcgacgcgcg ctaaagaatg gcaaaccaag 8820
tcgcgcgagc gtcgaaattc gtaatcatgg tcatagctgt ttcctgtgtg aaattgttat 8880
ccgctcaciaa ttccacacia catacgagcc ggaagcataa agtgtaaagc ctgggggtgcc 8940
taatgagtga gctaactcac attaatgtcg ttgcgctcac tgcccgcctt ccagtcggga 9000
aacctgtcgt gccagctgca ttaatgaatc ggccaacgcg cggggagagg cggtttgctg 9060
attgggcgct cttccgcttc ctgcctcact gactcgtgcg gctcggctcg tcggctgcgg 9120
cgagcgggtat cagctcactc aaaggcggta atacggttat ccacagaatc aggggataac 9180
gcaggaaaga acatgtgagc aaaaggccag caaaaggcca ggaaccgtaa aaaggccgcg 9240
ttgctggcgt ttttccatag gctccgcccc cctgacgagc atcacaaaaa tcgacgtca 9300
agtcagaggt ggcgaaaccc gacaggacta taaagatacc aggcgtttcc ccttgaagc 9360
tccctcgtgc gctctcctgt tccgacctg ccgcttaccg gatacctgtc cgcctttctc 9420
ccttcgggaa gcgtggcgct ttctcaatgc tcacgctgta ggtatctcag ttcgggtgtag 9480
gtcgttcgct ccaagctggg ctgtgtgcac gaacccccg ttcagcccg cgcgtgcgc 9540
ttatccggtg actatcgtct tgagtccaac ccggtgaagc acgacttatc gccactggca 9600
gcagccactg gtaacaggat tagcagagcg aggtatgtag gcggtgctac agagtcttg 9660
aagtgggtggc ctaactacgg ctacactaga aggacagtat ttggtatctg cgctctgctg 9720
aagccagtta ccttcggaaa aagagttggg agctcttgat ccggcaaaca aaccaccgct 9780
ggtagcgggt gtttttttgt ttgcaagcag cagattacgc gcagaaaaaa aggatctcaa 9840
gaagatecct tgatcttttc tacgggggtct gacgctcagt ggaacgaaaa ctcacgttaa 9900
gggatttttg tcatgagatt atcaaaaagg atcttcacct agatcctttt aaattaaaaa 9960
tgaagtttta aatcaatcta aagtatatat gagttaaact ggtctgacag ttaccaatgc 10020
ttaatcagtg aggcacctat ctacagcgtc tgtctatttc gttcatccat agttgcctga 10080
ctccccgtcg ttagataaac tacgatacgg gagggcttac catctggccc cagtgcctga 10140
atgataccgc gagaccacg ctacccggct ccagatttat cagcaataaa ccagccagcc 10200
ggaagggccg agcgcagaag tggctctgca actttatccg cctccatcca gtctattaat 10260
tggtgcccgg aagctagagt aagtagttcg ccagttaata gtttgcgcaa cgttgttgcc 10320
attgctacag gcatcgtggg gtcacgctcg tcgtttggta tggcttcatt cagctccggt 10380
tcccaacgat caaggcgagt tacatgatcc cccatgttgt gcaaaaaagc ggtagctcc 10440
ttcggctctc cgatcgttgt cagaagtaag ttggccgcag tggtatcact catgggtatg 10500
gcagcactgc ataattctct tactgtcatg ccatccgtaa gatgcttttc tgtgactggg 10560
gagtactcaa ccaagtcatt ctgagaatag tgtatgcggc gaccgagttg ctcttgcccc 10620
gcgtcaatac gggataatac cgcgccacat agcagaactt taaaagtgt catcattgga 10680
aaacgttctt cggggcgaaa actctcaagg atcttaccgc tggtgagatc cagttcgatg 10740
taaccactc gtgcacccaa ctgatcttca gcatctttta ctttcaccag cgtttctggg 10800

tgagcaaaaa caggaaggca aaatgccgca aaaaagggaa taagggcgac acggaaatgt 10860
 tgaataactca tactcttcct ttttcaatat tattgaagca tttatcaggg ttattgtctc 10920
 atgagcggat acatatattga atgtatttag aaaaataaac aaataggggt tccgcgcaca 10980
 tttccccgaa aagtgccacc tgacgtctaa gaaaccatta ttatcatgac attaacctat 11040
 aaaaataggc gtatcacgag gccctttcgt ctgcgcggtt tcggtgatga cggtgaaaac 11100
 ctctgacaca tgcagctccc ggagacgggtc acagcttggtc tgtaagcgga tgccgggagc 11160
 agacaagccc gtcagggcgc gtcagcgggt gttggcgggt gtcggggctg gcttaactat 11220
 gcggcatcag agcagattgt actgagagtg caccatatat gcggtgtgaa ataccgcaca 11280
 gatgcgtaag gagaaaatac cgcatcaggc gccattcgcc attcaggctg cgcaactggt 11340
 gggaagggcg atcgggtcgg gcctcttcgc tattacgcca gctggcgaaa gggggatgtg 11400
 ctgcaaggcg attaagttgg gtaacgccag ggttttccca gtcacgacgt tgtaaaacga 11460
 cgccagtgcc caagctttgt ttaaaatata aaaaattgt gatccacaa aatgaagtgg 11520
 ggcaaatca aataattaat agtgtccgta aacttggttg tcttcaactt tttgaggaac 11580
 acgttgagcg gcaaatccgt gactataaca caagttgatt taataatatt agccaacacg 11640
 tcgggctgcg tgttttttgc cgacgcgtct gtgtacacgt tgattaactg gtcgattaaa 11700
 ctgttgaaat aatttaattt ttggttcttc tttaaatctg tgatgaaatt ttttaaaata 11760
 actttaaat cttcattggt aaaaaatgcc acgttttgca acttggtgagg gtctaatatg 11820
 aggtcaaact cagtaggagt tttatccaaa aaagaaaaca tgattacgtc tgtacacgaa 11880
 cgcgtattaa cgcagagtgc aaagtataag agggttaaaa aatatatttt acgcaccata 11940
 tacgcatcgg gttgatatcg ttaatatgga tcaatttgaa cagttgatta acgtgtctct 12000
 gctcaagtct ttgatcaaaa cgcaaatcga cgaaaatgtg tcggacaata tcaagtcgat 12060
 gagcgaaaaa ctaaaaaggc tagaatacga caatctcaca gacagcgttg agatatacgg 12120
 tattcacgac agcaggctga ataataaaaa aattagaaac tattatttaa ccctagaaag 12180
 ataatacatat tgtgacgtac gttaaagata atcatgcgta aaattgacgc atgtgtttta 12240
 tcggtctgta tatcgagggt tatttattaa tttgaataga tattaagttt tattatattt 12300
 acacttacat actaataata aattcaacaa acaatttatt tatgtttatt tatttattaa 12360
 aaaaaaacia aaactcaaaa tttcttctat aaagtaacaa aacttttaaa cattctctct 12420
 tttacaaaaa taaacttatt ttgtacttta aaaacagtca tgttgtatta taaaataagt 12480
 aattagctta acttatacat aatagaaaca aattatactt attagtcagt cagaaacaac 12540
 tttggcacat atcaatatta tgctctcgac aaataacttt tttgcatttt ttgcacgatg 12600
 catttgccct tcgccttatt ttagaggggc agtaagtaca gtaagtacgt tttttcatta 12660
 ctggctcttc agtactgtca tctgatgtac caggcacttc atttggcaaa atattagaga 12720
 tattatcgcg caaatatctc ttcaaagtag gagcttctaa acgcttacgc ataaacgatg 12780
 acgtcaggct catgtaaagg tttctcataa attttttgcg actttggacc ttttctccct 12840

tgctactgac attatggctg tatataataa aagaatttat gcaggcaatg tttatcattc 12900
 cgtacaataa tgccataggc cacctattcg tcttctact gcaggtcac acagaacaca 12960
 tttgggtctag cgtgtccact cgccttttag tttgattata atacataacc atttgcggtt 13020
 taccgggagc ctcttaatta agggccccc tgcagaccat ggtcgatcga caaacacgaa 13080
 aatacattcc tggacagacc taggggtttg ctcttttttc tttgcaggat ctggattcaa 13140
 caaagaagga aaaatctacg gagtttggaa cctcatttat tgcacgaaga aacttagaga 13200
 gaaagactga tgcctaataa agactcaagc tgattgcaaa aacaatataa agttgtacct 13260
 tattgcggtg tcatcgtaaa accaagatgc ttaaaaggga ttagggccac ttaaattact 13320
 gcgaaaaatt ttattttacaa atttctcaca tgacttcag ataaactaat taactttgtg 13380
 tgtcaagggt gcttttccgg cagaaaagtc aatttgcaag tgatttaggt ttttttaatg 13440
 actatctttg caattgaatt gggtagtatt agatattgaa tataccaata ttgaggtata 13500
 atacaaatac aacaacattt gacctcggca actatcagaa gtacatgtag gtcgggaaca 13560
 ttagggaatt ctgcgcgata agttactaat tatgctaagc tatcgagaa cctcattaag 13620
 acttgagctg cagtgttttg caggtagtta tcatatttac agtattaatg tatgcagtat 13680
 tatattaatt ctaaataaga aagtcaggag aaggatattc aaaactgaag aattaatgta 13740
 cattagtttt gtttattaga ttttcaatat tgcagcatag aatatgtgta agtagtgag 13800
 agctaaacgc gcttttcaga acgttcatgc taattgctca tggatctcac aggaacaggt 13860
 ggtggcggcc ctccgtgcgc tcgtactgct ccacgatggt gtagtcctcg ttgtgggagg 13920
 tgatgtccag cttggcgctc acgtagtagt agccgggcag ctgcacgggc ttcttggtcca 13980
 tgtagatgga cttgaactcc accaggtagt ggccgcccgc cttcagcttc agggccttgt 14040
 ggggtctcgc cttcagcacg ccgtcgcggg ggtacaggcg ctccgtggag gcctcccagc 14100
 ccattggtctt cttctgcac acggggccgt cggaggggaa gttcacgcc atgaacttca 14160
 ccttgtagat gaagcagccg tcctgcaggg aggagtcctg ggtcacgggc gccacgccgc 14220
 cgtcctcgaa gttcatcacg cgtccact tgaagccctc ggggaaggac agcttcttgt 14280
 agtcggggat gtcggcgggg tgcttcacgt acaccttga gccgtactgg aactgggggg 14340
 acaggatgtc ccaggcgaag ggcagggggc cgccttgggt caccttcagc ttcacggtgt 14400
 tgtggccctc gtaggggagg ccctcgccct cgcctcgat ctgaaactcg tggccgttca 14460
 cgggtgccctc catgcgcacc ttgaagcgca tgaactcggg gatgacgttc tcggaggagg 14520
 ccattctagc ctttttggtt ctagatggca gttgtttatc tggaaaggag atggaagtgc 14580
 actcggtttag ctcgatctgt cgattatata tgtaaagttt tatagctccc acctacatct 14640
 caattcttta aattctacta ctttaatgac aacatagaaa actaatttgt gtctcaagaa 14700
 aatatttaac tttttaagat attgaagaag tgaagtgaat ctctattgtc catttgagca 14760
 catttcagta aataaaacca tgaaaaattg ccaaaattgg tatattccat ttccttttga 14820
 gtcattgctaa ttggttatct acttgaaact aaatcagtgc tctaaaaata tcggaaaaga 14880

```

atagttcata gaacttaatt taaacgtatc aaattaattg aaaatacgtt tcgttgcaat 14940
tactcttcac tttatatatta gatattgaac tttcgaagac cgattagtga cttgtagaaa 15000
aatccatcaa acaatccgaa cttattaaaa agagttttta tatccttagt cctcttggt 15060
ttttctgtaa taatccttgg atccttcgac attgagggtcg cactcacaca ccttttaggt 15120
caacgaaagg agaaaaattg gatagagtcg cacgttcgat tcctgttgac aactagattc 15180
aacaacaaat ggcacgccga tcggtctgtc ctgcctttat atctcggtc tccctcgaag 15240
cgatttgagt tggatgccag tgccggcactg agaaaaagat tgagcgaaag tgctgcggaa 15300
caatcaactc gaatcgatcc atataagcag ttttcagact tctggccgag catgtcagaa 15360
gtttggatgc ctcatcgac gaagacgact cattaatgcg cttcacaaaa aaaggaagcc 15420
acacaattaa atacttaata aagcatagaa tcatttatta agattaccaa taaattcaat 15480
aaacgaggga atttccaaac tgcgaaaata tgacatttta ttttttttc acattgttag 15540
taggtacaaa tcctagaaaa tgggtctcata acatataaca gcttaaataa ctatgtagta 15600
gaagataaac tttatgttca attcagttca tttttaatcc tagaaatatt taattaattg 15660
acaatatatg atggaagtaa gctatcaatc ttaggacctt attcattatt tgtagggagt 15720
gccaggagag agaaagctct cgaaagataa gagaacgcca tagccgagtg attagcagag 15780
gtgcgtgata gtgcgtgtat gggccacctt aacggcaatc agcactctcg agtggccaat 15840
tagcccatc tccgtcccca tctctgtccc aatcctgtcg ccaatgggcg actgctgcgg 15900
cggacgactc attcgggtgt gagagtccgg aaaatagaaa cggaccttag aaggaccgac 15960
gcaagacaat tggcatccgc ggcagggact gggagaatgg gacatttcaa cgggcacgca 16020
gcgaggcgaa tctaaaatag acgcgccaag cagagcggca cctataaatc aacatcaagg 16080
ttgatcgaga tatccacca tctgcggccg ccattgcgtca attttacgca gactatcttt 16140
ctagggttaa tcagccc 16157

```

<210> 4
 <211> 16157
 <212> DNA
 <213> artificial

<220>
 <223> predicted sequence of pLA1125 construct

```

<400> 4
gggcaattcg ggcgcgcctc ttaattaagg cccccctcg agaccatggt cgatcgacaa 60
acacgaaaat acattcctgg acagacctag ggggttgctc ttttttcttt gcaggatctg 120
gattcaacaa agaaggaaaa atctacggag tttggaacct catttattgc acgaagaaac 180
ttagagagaa agactgatgc ctaataaaga ctcaagctga ttgcaaaaac aatataaagt 240
tgtaccttat tgcggtgtca tcgtaaaacc aagatgctta aaagggatta ggccactta 300
aattactgcg aaaaatttta tttacaaatt tctcacatga cttccagata aactaattaa 360
ctttgtgtgt caaggttgct tttccggcag aaaagtcaat ttgcaagtga tttagggttt 420

```

tttaatgact	atctttgcaa	ttgaattggg	tactattaga	tattgaatat	accaatattg	480
aggataata	caaatacaac	aacatttgac	ctcggcaact	atcagaagta	catgtaggtc	540
gggaacatta	gggaattctg	cgcgataagt	tactaattat	gctaagctat	cgcagaacct	600
cattaagact	tgagctgcag	tgttttgcag	gtacttatca	tatttacagt	attaatgtat	660
gcagtattat	attaattcta	aataagaaa	tcaggagaag	gatattcaaa	actgaagaat	720
taatgtacat	tagttttggt	tattagattt	tcaatattgc	agcatagaat	atgtgtaagt	780
agtgagagc	taaacgcgct	tttcagaacg	ttcatgctaa	ttgctcatgg	atctcacagg	840
aacaggtggt	ggcggccctc	ggtgcgctcg	tactgctcca	cgatggtgta	gtcctcgttg	900
tgggaggtga	tgctccagctt	ggcgtccacg	tagtagtagc	cgggcagctg	cacgggcttc	960
ttggccatgt	agatggactt	gaactccacc	aggtagtggc	cgccgtcctt	cagcttcagg	1020
gccttggtgg	tctcgccctt	cagcacgccg	tcgcgggggt	acaggcgctc	ggtggaggcc	1080
tcccagccca	tggtcttctt	ctgcatcacg	gggccgtcgg	aggggaagt	cacgccgatg	1140
aacttcacct	tgtagatgaa	gcagccgtcc	tcagggagg	agtcctgggt	cacggctgcc	1200
acgccgccgt	cctcgaagtt	catcacgcgc	tcccacttga	agccctcggg	gaaggacagc	1260
ttcttgtagt	cggggatgtc	ggcgggggtc	ttcacgtaca	ccttgaggcc	gtactggaac	1320
tggggggaca	ggatgtccca	ggcgaagggc	agggggccgc	ccttggtcac	cttcagcttc	1380
acggtgttgt	ggccctcgta	ggggcgcccc	tcgccctcgc	cctcgatctc	gaactcgtgg	1440
ccgttcacgg	tgccctccat	gcgcaccttg	aagcgcatga	actcggtgat	gacgttctcg	1500
gaggaggcca	ttctagccat	tttggttcta	gatccgggggt	ctctggatta	gacgactgct	1560
ggctgatgga	gcggctttgt	gtcgggagga	gtatccacac	agcacaagaa	ctcaaacggt	1620
agtgatatga	aaactgggtc	cgcagcccggt	tttataaagc	ccgcttagcg	ctaaaacctt	1680
atcagccgca	atgctctccg	ctctcaagtc	gcgttcaaaa	cttttaccat	atatgggtat	1740
gaattgttcg	ccgattgggt	tttcatttac	ggagaatttc	ctccgcaact	gggtgttcgt	1800
gcttggtgtg	gtcttggtata	tcgccatgaa	gagtgtactt	ccatgggtcc	tatcagttgc	1860
ggcatgtgat	tcaatttggg	ggttactcat	attcgatact	tcagctggtt	atctttatgc	1920
tgtgatggtc	acggtgcttt	tacacatttt	ttccccaact	actcattgta	tgccaaaaca	1980
tacatatgta	catgtgttta	tagtgggttt	attagtaatg	tccagcgata	ttaaacatt	2040
acagttaaat	aatattcata	ttagcttacg	tttactgcta	tctgcacttc	aaattgggta	2100
ttcagggaca	tttggtgcaac	acacatcttc	tagtaggggc	atcactagtc	gtagaaaaca	2160
cacctaaact	aagccgggtt	taacaataac	ggcgaacag	ctgccagacg	agagaaaagt	2220
acacagcaag	aagcatttct	ataatgatgc	cgtcctaaat	atgcagcatt	tttgctcaaa	2280
ttgatcattt	cgttgcatcg	ctcaggtggt	atagtatttg	gctattgaat	gtgattgcta	2340
tcttaatagt	tttgaatagt	aatagtgcaa	atattcattc	ataaaaaata	ggtaatcgat	2400
tttctcagtg	cagtctatgc	gtttgcgcaa	acaggagacg	aagagagaga	cgcctcacac	2460

acaagcacat gcgtactacg ctggcctact ccgctactct atgtgtcgtc ctctgcatcg 2520
 tctgcttggc ggcagttctt ccagaggctg tgacgtcagg taagctatct cgtctcgggtg 2580
 tgctgttagc ccgtaacgat aagctgtccg attcctaatt ctctttcaaa taatttggtt 2640
 tatttttttt tttgtgtgtt ttgaagtcaa acagcttcat aaacaatatt aaatatcgca 2700
 attgttattg caaatgcatt cgtatcaggc gccttgatgt ttttcggcca aaacttcgtc 2760
 tgctctttgt cagcgttaatt gcgtttgccg gtcgtgccgt cttgctcgca cgcgcttgca 2820
 aacaaaaagt gaaagaggga aagccactgt gttgttgctt ttgttgctat cgttggtgctc 2880
 acacacatgg caataagtaa ccatattgta gacttattta ttttagcgtt ttcgttgact 2940
 aatctttgtt taaataattg cctatttcga gccaaaatga cgcataaagg acacggcaac 3000
 atgcggtgctg ttgagcatgc gtacgagatg gcgctagaga gagcgagata aaaagaaaaa 3060
 aacaaccggc gagtggcgag acacgaataa tacacacacg cacacatggt caaaggagtg 3120
 gctccaacgt cggtgacgtc gcgcccagcc acagaccccg ccacgccccg ccccaccctt 3180
 ccttcgctg ccccgacccc ctttccttcc actaccgttt tgccgcattt ggtattgcca 3240
 tctcgtcttt ttggctttac tttgcttttt gaataagcca agaaaactgaa atcaaagcgg 3300
 acagtccgaa aattgaaact attccaaaaa acaacaaccg cacatgcact gcgttcggca 3360
 aaagacgcag aaatcagcct gtggtctggc atcatcttaa aatatcatca gtatcatcat 3420
 tgccattata aatacataag aaaatgagaa tacattaaag gtatatacat acaatctttg 3480
 tcatagtcaa catattttgc aacaaaatta tctatagata acatgttttt gtagaaaatt 3540
 ccactaactg attttcgtat aaatacaaat tattctcatt cttcaaaaata ctcgccgacg 3600
 tcatcgcgaa attcctcagc attcttcgac aaatgaagag agacagcgag tgacacaatc 3660
 ttatcttacc gacgagtctt tgaaaatggc cccgggaacg gtgaaaaaca tatagccaag 3720
 aatggagcga tcgccaggcg atggaaattg gcttcaaatt gctccgaata tggccagttg 3780
 gtcagttggc attggattgt ggagagatac tagatcagct cgttctcctt ctctccgca 3840
 ttctcctact cgcccgaaga cgcttctttg atcgcttctt tctcgaatgt acatattgat 3900
 atgtatttgg tgtttggcga cgatcagcca gctttaatgg ccataaaacg tgtcttggtt 3960
 tgtggggccac ataacaggca gcgagatctg gaaaaatgat gtgacagtgg aaatgagaag 4020
 taggtgcac tgcaaagatt tgattcagag ttgatgccat tcatgatctt ttttagaagt 4080
 atcatttgtg tttttagaat atagaattct agtatgtatg taagttaata aaaccattt 4140
 ttgcggaaag tagataaaaa aaacattttt tttttttact gcactggata tctgcggccg 4200
 gccagttaa ccctagaaag ataatcatat tgtgacgtac gttaaagata atcatgcgta 4260
 aaattgacgc atgggtacca gatatccagt gcagtaaaaa aaaaaaatgt tttttttatc 4320
 tactttccgc aaaaatgggt tttattaact tacatacata ctagaattct aaaaaaatc 4380
 atgaatggca tcaactctga atcaaatctt tgcagatgca cctacttctc atttccactg 4440
 tcacatcatt tttccagatc tcgctgcctg ttatgtggcc cacaaccaa gacacgtttt 4500

atggccatta aagctggctg atcgtcgcca aacaccaa	at	at	at	at	at	4560
gagaaagaag cgatcaaaga agcgtcttcg ggcgagtagg	aga	at	gc	ga	gg	4620
aacgagctga tctagtatct ctccacaatc caatgccaac	tg	acca	actg	gcc	atattcg	4680
gagcaatttg aagccaattt ccacgcctg gcgatcgctc	catt	ctctggc	tata	tgtttt		4740
tcaccgttcc cggggccatt ttcaaagact cgtcggtg	aa	ata	agattgt	gtc	actcgct	4800
gtctctcttc atttgtcgaa gaatgctgag gaatttcg	cg	at	gcgtcg	cg	agtatttt	4860
gaagaatgag aataatttgt atttatacga aaatcagt	ta	gt	gaatttt	ct	acaaaaac	4920
atgttatcta tagataattt tgttgcaaaa tatgttgact	at	gc	aaa	ga	ttgtatgtat	4980
atacctttaa tgtattctca ttttcttatg tatttataat	gg	caatgatg	ata	ctgatga		5040
tattttaaga tgatgccaga ccacaggctg atttctg	cg	cttttgc	ga	acgcagtga		5100
tgtgcggttg ttgttttttg gaatagtttc aattttcg	ga	ctgtccg	ctt	tgatttcagt		5160
ttcttggtt attcaaaaag caaagtaaag ccaaaaag	gc	gagatgg	caa	tac	caa	5220
ggcaaaacgg tagtggaagg aaaggggtgc ggggcag	cg	aag	gaagggt	ggggc	ggggc	5280
gtggcggttg ctgtggctgg gcgcgacgtc accgac	gtt	gagccactcc	ttt	gaccatg		5340
tgtgcgtgtg tgtattattc gtgtctcgcc actcg	cggt	tg	tttttttc	tttttatctc		5400
gctctctcta ggcctatctc gtacgcattc tcaacgc	acc	gc	atgttgcc	gtgtccttta		5460
tgcgtcattt tggctcgaaa taggcaatta tttaaaca	aa	gattag	tcaa	cgaaaacgct		5520
aaaataaata agtctacaat atgggttactt attgcc	atgt	gtgtgcagcc	aac	gatagca		5580
acaaaagcaa caacacagtg gctttccctc tttcact	ttt	tg	tttgcaag	cg	cg	5640
caagacggca cgaccggcaa acgcaattac gctgaca	aa	agc	gacgaa	gttttgccg		5700
aaaaacatca aggcgcctga tacgaatgca tttgca	ata	ca	attg	cgat	attta	5760
gtttatgaag ctgtttgact tcaaaacaca caaaaaa	aa	aata	aa	caa	attattg	5820
agagaattag gaatcgga gcttatcggt acgggcta	ac	agc	acacga	gac	gaa	5880
cttacctgac gtcacagcct ctggaagaac tgccgca	ag	cag	acgatgc	agaggacgac		5940
acatagagta ggcgagtagg ccagcgtagt acgcat	gtgc	ttgtgtgtga	ggcgtctctc			6000
tcttcgtctc ctgtttgcgc aaacgcatag actgc	actga	gaaaatcgat	tac	ctatttt		6060
ttatgaatga atatttgac tattactatt caaaact	att	aagatag	caa	tcacatt	caa	6120
tagccaaata ctataccacc tgagcgatgc aacgaa	tga	tcaatttgag	caaaa	atgct		6180
gcatatttag gacggcatca ttatagaaat gcttct	tgct	gtgtactttt	ctctcgctg			6240
gcagctgttt cgcggttatt gttaaaaccg gctta	ag	tta	ggtgtgtttt	ctacgactag		6300
tgatgccct actagaagat gtgtgttgca caaatg	tccc	tgaataacca	atttgaagt			6360
cagatagcag taaacgtaag ctaatatgaa tattatt	taa	ctg	taatgtt	ttaatatgc		6420
tggacattac taataaaccc actataaaca catgt	acata	tgtatgtttt	ggcatacaat			6480
gagtagttgg ggaaaaatg tgtaaaagca ccgtg	accat	cacagcataa	agataaccag			6540

ctgaagtatc gaatatgagt aacccccaaa ttgaatcaca tgccgcaact gataggaccc	6600
atggaagtac actcttcatg gcgatataca agacacacac aagcacgaac acccagttgc	6660
ggaggaaatt ctccgtaaatt gaaaacccaa tcggcggaaca attcataccc atatatggta	6720
aaagttttga acgcgacttg agagcggaga gcattgcggc tgataagggt ttagcgctaa	6780
gcgggcttta taaaacgggc tgcgggacca gttttcatat cactaccgtt tgagttcttg	6840
tgctgtgtgg atactcctcc cgacacaaaag ccgctccatc agccagcagt cgtctaattc	6900
agagaccccg gatctagaac caaaatggct agcaaaggag aagaactttt cactggagtt	6960
gtcccaattc ttgttgaatt agatggtgat gttaatgggc acaaattttc tgtcagtga	7020
gaggggtgaag gtgatgcaac atacggaaaa cttaccctta aatttatttg cactactgga	7080
aaactacctg ttccatggcc aacacttgtc actactttca cttatggtgt tcaatgcttt	7140
tcaagatacc cagatcatat gaaacagcat gactttttca agagtgccat gccgaagggt	7200
tatgtacagg aaagaactat atttttcaaa gatgacggga actacaagac acgtgctgaa	7260
gtcaagtttg aagggtgatac cttgttaat agaatcgagt taaaagggtat tgattttaaa	7320
gaagatggaa acattcttgg acacaaattg gaatacaact ataactcaca caatgtatac	7380
atcatggcag acaaacaaaa gaatggaatc aaagttaact tcaaaattag acacaacatt	7440
gaagatggaa gcgttcaact agcagaccat tatcaacaaa atactccaat tggcgatggc	7500
cctgtccttt taccagacaa ccattacctg tccacacaat ctgccctttc gaaagatccc	7560
aacgaaaaga gagaccacat ggtccttctt gagtttgtaa cagctgctgg gattacacat	7620
ggcatggatg aactatacaa ataaggatcc atgagcaatt agcatgaacg ttctgaaaag	7680
cgcgtttagc tctccactac ttacacatat tctatgctgc aatattgaaa atctaataaa	7740
caaaactaat gtacattaat tcttcagttt tgaatatcct tctcctgact ttcttattta	7800
gaattaatat aatactgcat acattaatac tgtaaatatg ataagtacct gcaaaacact	7860
gcagctcaag tcttaatgag gttctgcgat agcttagcat aattagtaac ttatcgcgca	7920
gaattcccta atgttcccga cctacatgta cttctgatag ttgccgaggt caaatgttgt	7980
tgtatttgta ttatacctca atattggtat attcaatata taatagtacc caattcaatt	8040
gcaaagatag tcattaaaaa aacctaaatc acttgcaaat tgacttttct gccggaaaag	8100
caaccttgac acacaaagtt aattagttta tctggaagtc atgtgagaaa tttgtaaata	8160
aaatttttcg cagtaattta agtgggccta atccctttta agcatcttgg ttttacgatg	8220
acaccgcaat aagggtacaac tttatattgt ttttgcaatc agcttgagtc tttattaggc	8280
atcagtcttt ctctctaagt ttcttcgtgc aataaatgag gttccaaaact ccgtagattt	8340
ttccttcttt gttgaatcca gatcctgcaa agaaaaaaga gcaaaccctt aggtctgtcc	8400
aggaatgtat tttcgtgttt gtcgatcgac catggtcacc ggtagatata tataacaaga	8460
aaatatatat ataataagtt atcacgtaag tagaacatga aataacaata taattatcgt	8520
atgagttaaa tcttaaaagt cacgtaaaag ataatcatgc gtcattttga ctcacgcggt	8580

cggtatagtt	caaaatcagt	gacacttacc	gcattgacaa	gcacgcctca	cgggagctcc	8640
aagcggcgac	tgagatgtcc	taaatgcaca	gcgacggatt	cgcgctatct	agaaagagag	8700
agcaatatct	caagaatgca	tgcgtaatt	ttacgcagac	tatctttcta	gggttaagtc	8760
gacatccaca	ttgtggccaa	gctgtgtgac	gcgacgcgcg	ctaaagaatg	gcaaaccaag	8820
tcgcgcgagc	gtcgaaattc	gtaatcatgg	tcatagctgt	ttcctgtgtg	aaattgttat	8880
ccgctcacia	ttccacacaa	catacgagcc	ggaagcataa	agtgtaaagc	ctgggggtgcc	8940
taatgagtga	gctaactcac	attaattgcg	ttgcgctcac	tgcccgcttt	ccagtcggga	9000
aacctgtcgt	gccagctgca	ttaatgaatc	ggccaacgcg	cggggagagg	cggtttgcgt	9060
attgggcgct	cttcgccttc	ctcgcctact	gactcgcctg	gctcggctcg	tcggctgcgg	9120
cgagcgggat	cagctcactc	aaaggcggta	atacggttat	ccacagaatc	aggggataac	9180
gcaggaaga	acatgtgagc	aaaaggccag	caaaaggcca	ggaaccgtaa	aaaggccgcg	9240
ttgctggcgt	ttttccatag	gtccgccttc	cctgacgagc	atcacaaaaa	tcgacgctca	9300
agtcagaggt	ggcgaaacct	gacaggacta	taaagatacc	aggcgcttcc	ccctggaagc	9360
tcctcgtgc	gctctcctgt	tcgacccctg	ccgcttaccc	gatacctgtc	cgcctttctc	9420
ccttcgggaa	gcgtggcgct	ttctcaatgc	tcacgctgta	ggtatctcag	ttcgggtgtag	9480
gtcgcttcgt	ccaagctggg	ctgtgtgcac	gaaccccccg	ttcagcccca	ccgctgcgcc	9540
ttatccggta	actatcgtct	tgagtccaac	ccggttaagc	acgacttata	gccactggca	9600
gcagccactg	gtaacaggat	tagcagagcg	aggtatgtag	gcggtgctac	agagttcttg	9660
aagtgggtgg	ctaactacgg	ctacactaga	aggacagtat	ttggtatctg	cgcctctgctg	9720
aagccagtta	ccttcggaaa	aagagttggg	agctcttgat	ccggcaaaaa	aaccaccgct	9780
ggtagcgggt	gtttttttgt	ttgcaagcag	cagattacgc	gcagaaaaaa	aggatctcaa	9840
gaagatcctt	tgatcttttc	tacgggggtc	gacgctcagt	ggaacgaaaa	ctcacgttaa	9900
gggatttttg	tcatgagatt	atcaaaaagg	atcttcacct	agatcctttt	aaattaaaaa	9960
tgaagtttta	aatcaatcta	aagtatatat	gagtaaactt	ggtctgacag	ttaccaatgc	10020
ttaatcagtg	aggcacctat	ctcagcgatc	tgtctatttc	gttcatccat	agttgcctga	10080
ctccccgtcg	tgtagataac	tacgatacgg	gagggtttac	catctggccc	cagtgcctga	10140
atgataccgc	gagacccacg	ctcaccggct	ccagatttat	cagcaataaa	ccagccagcc	10200
ggaagggccg	agcgcagaag	tggtcctgca	actttatccg	cctccatcca	gtctattaat	10260
tggttgcggg	aagctagagt	aagtagttcg	ccagtttaata	gtttgcgcaa	cggtgttgcc	10320
attgctacag	gcatcggtgt	gtcacgctcg	tcgtttggta	tggcttcatt	cagctccggg	10380
tcccaacgat	caaggcgagt	tacatgatcc	cccatgttgt	gcaaaaaagc	ggtagctcc	10440
ttcggctctc	cgatcgttgt	cagaagtaag	ttggccgcag	tggtatcact	catggttatg	10500
gcagcactgc	ataattctct	tactgtcatg	ccatccgtaa	gatgcttttc	tgtgactggg	10560
gagtactcaa	ccaagtcatt	ctgagaatag	tgtatgcggc	gaccgagttg	ctcttgcccc	10620

gcgtcaatac gggataatac cgcgccacat agcagaactt taaaagtgt catcattgga 10680
 aaacgttctt cggggcgaaa actctcaagg atcttaccgc tgttgagatc cagttcgatg 10740
 taaccactc gtgcacccaa ctgatcttca gcatctttta ctttcaccag cgtttctggg 10800
 tgagcaaaaa caggaaggca aaatgccgca aaaaaggga taagggcgac acggaaatgt 10860
 tgaatactca tactcttcct ttttcaatat tattgaagca tttatcaggg ttattgtctc 10920
 atgagcggat acatatattga atgtatttag aaaaataaac aaataggggt tccgcgcaca 10980
 tttccccgaa aagtgccacc tgacgtctaa gaaaccatta ttatcatgac attaacctat 11040
 aaaaataggc gtatcacgag gccctttcgt ctgcgcggtt tcggtgatga cggtgaaaac 11100
 ctctgacaca tgcagctccc ggagacgggt acagcttggt tgtaagcggg tgccgggagc 11160
 agacaagccc gtcagggcgc gtcagcgggt gttggcgggt gtcggggctg gcttaactat 11220
 gcggcatcag agcagattgt actgagagtg caccatatat gcggtgtgaa ataccgcaca 11280
 gatgcgtaag gagaaaatac cgcatacaggc gccattcgcc attcaggctg cgcaactgtt 11340
 gggaaggcg atcgggtcgg gcctcttcgc tattacgcca gctggcgaaa gggggatgtg 11400
 ctgcaaggcg attaatgttg gtaacgccag ggttttccca gtcacgacgt tgtaaaacga 11460
 cggccagtgc caagctttgt ttaaaatata aaaaaattgt gatcccacaa aatgaagtgg 11520
 ggcaaaatca aataattaat agtgtccgta aacttggttg tcttcaactt tttgaggaac 11580
 acgttgagc gcaaatccgt gactataaca caagttgatt taataatttt agccaacacg 11640
 tcgggctcgc tgttttttgc cgacgcgtct gtgtacacgt tgattaactg gtcgattaaa 11700
 ctgttgaaat aatttaattt ttggttcttc tttaaatctg tgatgaaatt ttttaaaata 11760
 actttaaatt cttcattggt aaaaaatgcc acgttttgca acttgtagg gtctaatatg 11820
 aggtcaaact cagtaggagt tttatccaaa aaagaaaaca tgattacgtc tgtacacgaa 11880
 cgcgatttaa cgcagagtgc aaagtataag agggttaaaa aatatatttt acgcaccata 11940
 tacgcacggt gttgatatcg ttaatatgga tcaatttgaa cagttgatta acgtgtctct 12000
 gctcaagtct ttgatcaaaa cgcaaatcga cgaaaatgtg tcggacaata tcaagtcgat 12060
 gagcgaaaaa ctaaaaaggc tagaatacga caatctcaca gacagcgttg agatatacgg 12120
 tattcacgac agcaggctga ataataaaaa aattagaaac tattatttaa ccctagaaag 12180
 ataatcatat tgtgacgtac gttaaagata atcatgcgta aaattgacgc atgtgtttta 12240
 tcggctctgta tatcgagggt tattttattaa ttgaaataga tattaagttt tattatattt 12300
 acattacat actaataata aattcaacaa acaatttatt tatgtttatt tttttattaa 12360
 aaaaaacaa aaactcaaaa tttcttctat aaagtaacaa aacttttaa cattctctct 12420
 tttacaaaaa taaacttatt ttgtacttta aaaacagtca tgttgattta taaaataagt 12480
 aattagctta acttatacat aatagaaaca aattatactt attagtcagt cagaaacaac 12540
 tttggcacat atcaatatta tgctctcgac aaataacttt tttgcatttt ttgcacgatg 12600
 catttgctt tcgcttatt ttagaggggc agtaagtaca gtaagtacgt tttttcatta 12660

ctggctcttc agtactgtca tctgatgtac caggcacttc atttggcaaa atattagaga 12720
tattatcgcg caaatatctc ttcaaagtag gagcttctaa acgcttacgc ataaacgatg 12780
acgtcagget catgtaaagg tttctcataa attttttgcg actttggacc tttctccct 12840
tgctactgac attatggctg tatataataa aagaatttat gcaggcaatg tttatcatc 12900
cgtacaataa tgccataggc cacctattcg tcttctact gcaggtcac acagaacaca 12960
tttggcttag cgtgtccact cgccttttag tttgattata atacataacc atttgcggtt 13020
taccgggcg cttttaatta agggcccccc tcgagaccat ggtcgatcga caaacacgaa 13080
aatacattcc tggacagacc taggggtttg ctcttttttc tttgcaggat ctggattcaa 13140
caaagaagga aaaatctacg gagtttggaa cctcatttat tgcacgaaga aacttagaga 13200
gaaagactga tgcctaataa agactcaagc tgattgcaaa aacaatataa agttgtacct 13260
tattgcggtg tcatcgtaaa accaagatgc ttaaaggga ttagggccac ttaaattact 13320
gcgaaaaatt ttatttacia atttctcaca tgacttccag ataaactaat taactttgtg 13380
tgtcaagggt gcttttccgg cagaaaagtc aatttgcaag tgatttaggt ttttttaatg 13440
actatctttg caattgaatt ggttactatt agatattgaa tataccaata ttgaggtata 13500
atacaaatac aacaacattt gacctcggca actatcagaa gtacatgtag gtcgggaaca 13560
ttagggaatt ctgcgcgata agttactaat tatgctaagc tatcgcagaa cctcattaag 13620
acttgagctg cagtgttttg caggactta tcatatttac agtattaatg tatgcagtat 13680
tatattaatt ctaaataaga aagtcaggag aaggatattc aaaactgaag aattaatgta 13740
cattagtttt gtttattaga ttttcaatat tgcagcatag aatatgtgta agtagtggag 13800
agctaaacgc gcttttcaga acgttcatgc taattgctca tggatctcac aggaacagg 13860
gggtggcgcc ctcggtgcgc tcgtactgct ccacgatggt gtagtcctcg ttgtgggagg 13920
tgatgtccag cttggcgctc acgtagtagt agccgggcag ctgcacgggc ttcttgcca 13980
tgtagatgga cttgaactcc accaggtagt ggccgccgtc cttcagcttc agggccttgt 14040
gggtctcgcc cttcagcacg ccgtcgcggg ggtacaggcg ctcggtggag gcctcccagc 14100
ccatggtctt cttctgcac acggggccgt cggaggggaa gttcacgccg atgaacttca 14160
ccttgtagat gaagcagccg tcctgcaggg aggagtctg ggtcacggtc gccacgccgc 14220
cgtcctcgaa gttcatcacg cgctcccact tgaagccctc ggggaaggac agcttcttgt 14280
agtcggggat gtcggcgggg tgcttcacgt acaccttga gccgtactgg aactgggggg 14340
acaggatgtc ccaggcgaag ggcagggggc cgcccttggc caccttcagc ttcacgggtg 14400
tgtggccctc gtagggggcg ccctcgccct cgccctcgat ctggaactcg tggccgttca 14460
cgggtgccct catgcgcacc ttgaagcgca tgaactcggg gatgacgttc tcggaggagg 14520
ccattctagc cattttggtt ctagatggca gttgtttatc tggaaggagg atggaagtgc 14580
actcgggttag ctcgatctgt cgattatata tgtaaagttt tatagctccc acctacatct 14640
caattcttta aattctacta ctttaatgac aacatagaaa actaatttgt gtctcaagaa 14700

aatattttaac tttttaagat attgaagaag tgaagtgaaa ctctattgtc catttgagca 14760
catttcagta aataaaacca tgaaaaattg ccaaaattgg tatattccat ttccttttga 14820
gtcatgctaa ttggttattc acttgaaact aaatcagtg ctaaaaaata tcggaaaaga 14880
atagttcata gaacttaatt taaacgtatc aaattaattg aaaatacgtt tcgttgcaat 14940
tactcttcac tttatattta gatattgaac tttcgaagac cgattagtga cttgtagaaa 15000
aatccatcaa acaatccgaa cttattaaaa agagttttaa tatccttagt cctcttggtc 15060
ttttctgtaa taatccttgg atccttcgac attgaggtcg cactcacaca ccttttaggt 15120
caacgaaagg agaaaaattg gatagagtcg cacgttcgat tcctgttgac aactagattc 15180
aacaacaaat ggcacgccga tcggtctgtc ctgcctttat atctcggctc tccctcgaag 15240
cgatttgagt tggatgccag tgcggcactg agaaaaagat tgagcgaaag tgctgcggaa 15300
caatcaactc gaatcgatcc atataagcag ttttcagact tctggccgag catgtcagaa 15360
gtttggatgc ctcatcgac gaagacgact cattaatgcg cttcacaaaa aaaggaagcc 15420
acacaattaa atacttaata aagcatagaa tcatttatta agattaccaa taaattcaat 15480
aaacgaggga atttccaaac tgcgaaaata tgacatttta tatttttttc acattgttag 15540
taggtacaaa tcctagaaaa tggcttcata acatataaca gcttaaataa ctatgtagta 15600
gaagataaac tttatgttca attcagttca tatttaatcc tagaaatatt taattaattg 15660
acaatatatg atggaagtaa gctatcaatc ttaggaccct attcattatt tgtagggagt 15720
gccaggagag agaaagctct cgaaagataa gagaacgcca tagccgagtg attagcagag 15780
gtgcgtgata gtcggtgtat gggccaccta aacggcaatc agcactctcg agtggccaat 15840
tagcccatc tccgtcccca tctctgtccc aatcctgctg ccaatgggag actgctgcgg 15900
cggacgactc attcgtgtct gagagtctcg aaaatagaaa cggaccttag aaggaccgac 15960
gcaagacaat tggcatccgc ggcagggact gggagaatgg gacatttcaa cgggcacgca 16020
gcgaggcgaa tctaaaatag acgcgccaag cagagcggca cctataaatc aacatcaagg 16080
ttgatcgaga tatccacca tctgcggccg ccatgcgtca attttacgca gactatcttt 16140
ctagggttaa tcagccc 16157

<210> 5
<211> 531
<212> DNA
<213> *Drosophila melanogaster*

<400> 5
gaaatcgatt gcggaacca ttgcttagtt tttgattgac ctttgtttga aaacaaaata 60
tatgtataaa ggaattattt gaaagtacga gcggtttgtt gtcattattg tgggattttt 120
ggctatttca ttttaattaa gttttttgaa tttaaatgtt tcatttagct acattcagaa 180
atattgtttt gaatttaaaa aatttacagt gtgaatgtct tgaatttttc aattggaaaa 240
aaaaggatga tggttgaagt atccgctttg tttattacag tagactggtc aactgccag 300
ctgctaaata atagaaaaca gtctgcgtgg acaaacattt aattaaaatg aatgtaagca 360

ctttttaacg aaatctttgg gaatatttcg ctcacagca ttttatttga gcaggagtcc 420
gagtcccat cgacatacac acgctgaagc tgcaggactg gctgatcagc cgcagaattg 480
tgccaagaat gtgcagcagg agcttaggga gatccacagg aagattagca a 531

<210> 6
<211> 25
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 6
gccagtcttg cagcttcagc gtgtg 25

<210> 7
<211> 26
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 7
aaaggtatga tgggtgaagt atccgc 26

<210> 8
<211> 26
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 8
ccttctttgt tgaatccaga tcctgc 26

<210> 9
<211> 23
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 9
gctcctgggc aacgtgctgg ttg 23

<210> 10
<211> 32
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 10
ggtggctagc ttgcgcttct tcttgggtgg gg 32

<210> 11
<211> 53
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 11
ggtgggcgcg cccaattgcc cgggcttttt aaccctagaa agatagtctg cgt 53

<210> 12
<211> 40
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 12
gcccaagctt gcggccgcgt cattttgact cacgcggctg 40

<210> 13
<211> 29
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 13
gatttgattc agagttgatg ccattcatg 29

<210> 14
<211> 25
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 14
ccagacctcc accttcaagg tgacc 25

<210> 15
<211> 27
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 15
ttgtaggagg tgtggaactg gcatctg 27

<210> 16
<211> 25
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 16
caacaccgtg aagctgaagg tgacc 25

<210> 17
<211> 28
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 17
cttggccatg tagatggact tgaactcc 28

<210> 18
<211> 22
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 18
caagcaggcc atcaacctgt gc 22

<210> 19
<211> 24
<212> DNA
<213> artificial

<220>
<223> PCR Primer

<400> 19
gacttggcct tgtacacggt gtcg 24

<210> 20
<211> 10
<212> DNA
<213> artificial sequence

<220>
<223> consensus sequence

<220>
<221> misc_feature
<222> (2)..(2)
<223> n is a, c, g, or t

<220>
<221> misc_feature
<222> (9)..(9)
<223> n is a, c, g, or t

<400> 20
wnwttawnw 10